

# AFTERGLOW OBSERVATIONS OF *Fermi*-LAT GAMMA-RAY BURSTS AND THE EMERGING CLASS OF HYPER-ENERGETIC EVENTS

S. B. CENKO<sup>1</sup>, D. A. FRAIL<sup>2</sup>, F. A. HARRISON<sup>3</sup>, J. B. HAISLIP<sup>4</sup>, D. E. REICHART<sup>4</sup>, N. R. BUTLER<sup>1,5</sup>, B. E. COBB<sup>1</sup>,  
A. CUCCHIARA<sup>6</sup>, E. BERGER<sup>7</sup>, J. S. BLOOM<sup>1</sup>, P. CHANDRA<sup>8</sup>, D. B. FOX<sup>6</sup>, D. A. PERLEY<sup>1</sup>, J. X. PROCHASKA<sup>9</sup>,  
A. V. FILIPPENKO<sup>1</sup>, K. GLAZEBROOK<sup>10</sup>, K. M. IVARSEN<sup>4</sup>, M. M. KASLIWAL<sup>11</sup>, S. R. KULKARNI<sup>11</sup>, A. P. LACLUYZE<sup>4</sup>,  
S. LOPEZ<sup>12</sup>, A. N. MORGAN<sup>1</sup>, M. PETTINI<sup>13,14</sup>, AND V. R. RANA<sup>3</sup>

*Submitted to ApJ*

## ABSTRACT

We present broadband (radio, optical, and X-ray) light curves and spectra of the afterglows of four long-duration gamma-ray bursts (GRBs 090323, 090328, 090902B, and 090926A) detected by the Gamma-Ray Burst Monitor (GBM) and Large Area Telescope (LAT) instruments on the *Fermi* satellite. With its widest spectral bandpass, extending to GeV energies, *Fermi* is sensitive to GRBs with very large isotropic energy releases ( $10^{54}$  erg). Although rare, these events are particularly important for testing GRB central-engine models. When combined with spectroscopic redshifts, our afterglow data for these four events are able to constrain jet collimation angles, the density structure of the circumburst medium, and both the true radiated energy release and the kinetic energy of the outflows. In agreement with our earlier work, we find that the relativistic energy budget of at least one of these events (GRB 090926A) exceeds the canonical value of  $10^{51}$  erg by an order of magnitude. Such energies pose a severe challenge for models in which the GRB is powered by a magnetar or neutrino-driven collapsar, but remain compatible with theoretical expectations for magneto-hydrodynamical collapsar models (e.g., the Blandford-Znajek mechanism). Our jet opening angles ( $\theta$ ) are similar to those found for pre-*Fermi* GRBs, but the large initial Lorentz factors ( $\Gamma_0$ ) inferred from the detection of GeV photons imply  $\theta\Gamma_0 \approx 70$ –90, values which are above those predicted in magnetohydrodynamic models of jet acceleration. Finally, we find that these *Fermi*-LAT events preferentially occur in a low-density circumburst environment, and we speculate that this might result from the lower mass-loss rates of their lower-metallicity progenitor stars. Future studies of *Fermi*-LAT afterglows in the radio with the order-of-magnitude improvement in sensitivity offered by the Extended Very Large Array should definitively establish the relativistic energy budgets of these events.

*Subject headings:* cosmology: observations - gamma rays: bursts - radio continuum: general

## 1. INTRODUCTION

<sup>1</sup> Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA.

<sup>2</sup> National Radio Astronomy Observatory, 1003 Lopezville Road, Socorro, NM 87801, USA.

<sup>3</sup> Space Radiation Laboratory, California Institute of Technology, MS 105-24, Pasadena, CA 91125, USA.

<sup>4</sup> Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC 27599, USA.

<sup>5</sup> Einstein Fellow.

<sup>6</sup> Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802, USA.

<sup>7</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA.

<sup>8</sup> Department of Physics, Royal Military College of Canada, Kingston, ON, Canada.

<sup>9</sup> Department of Astronomy and Astrophysics, UCO/Lick Observatory, University of California, 1156 High Street, Santa Cruz, CA 95064, USA.

<sup>10</sup> Centre for Astrophysics and Supercomputing, Swinburne University of Technology, 1 Alfred St, Hawthorn, Victoria 3122, Australia.

<sup>11</sup> Department of Astronomy, California Institute of Technology, MS 105-24, Pasadena, CA 91125, USA.

<sup>12</sup> Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile.

<sup>13</sup> Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, UK.

<sup>14</sup> International Center for Radio Astronomy Research, University of Western Australia, 35 Stirling Hwy, Crawley, WA 6009, Australia.

Long-duration gamma-ray bursts (GRBs<sup>15</sup>), like hydrogen-deficient Type Ib/c supernovae (SNe Ib/c), result from the gravitational collapse of the evolved core of a massive star. The main characteristic that sets GRBs apart from other SNe is that a substantial fraction of the energy of the explosion is coupled to relativistic ejecta. A compact central engine is responsible for accelerating and collimating these jet-like outflows and driving the SN explosions (Woosley & Bloom 2006; Gehrels et al. 2009; Soderberg et al. 2010). The precise nature of the central engine which powers GRB-SNe, however, remains an open question.

Motivated by empirical constraints, all viable central-engine models for long-duration GRBs share some common characteristics (e.g., Piran 2005). They must produce a collimated outflow with an initial Lorentz factor ( $\Gamma_0$ ) of a few hundred on observed time scales of 10–100 s, with luminosities and kinetic energies of order  $10^{50}$  erg s<sup>−1</sup> and  $10^{51}$  erg, respectively. Leading models include the “collapsar” model in which a relativistic jet is produced from a rotating black hole/accretion disk system (Woosley 1993; MacFadyen & Woosley 1999), and the “magnetar” model in which the rapid energy loss from a newly born millisecond neutron star (formed either from

<sup>15</sup> Throughout this work, we shall refer exclusively to long-duration GRBs (i.e., those apparently with massive-star progenitors) unless explicitly stated otherwise.

the gravitational collapse of a massive star or from accreting or coalescing white dwarfs) with a  $10^{15}$  G magnetic field drives a Poynting flux-dominated relativistic outflow (Usov 1992).

These and other more exotic models for GRB central engines are highly constrained by their energetics. The prompt high-energy emission, when combined with a spectroscopically determined redshift (and hence distance measurement), yields the isotropic radiated gamma-ray energy ( $E_{\gamma,\text{iso}}$ ). Well-sampled afterglow observations allow both a measurement of the degree of collimation (and hence the true beaming-corrected energy release in the prompt emission,  $E_{\gamma}$ ) and the kinetic energy remaining in the shock that powers the broadband afterglow emission ( $E_{\text{KE}}$ ). Such measurements, made nearly a decade ago, pointed to a total *relativistic* energy yield ( $E_{\text{rel}} \approx E_{\gamma} + E_{\text{KE}}$ ) of  $\sim 10^{51}$  erg (Frail et al. 2001; Panaitescu & Kumar 2001a; Freedman & Waxman 2001; Bloom et al. 2003; Berger et al. 2003a).

Since that time, there has been growing evidence for a considerable range in the relativistic energy scale  $E_{\text{rel}}$ , suggesting either a diversity in central engines or their properties. Most notably, a population of nearby (redshift  $z \lesssim 0.1$ ) subenergetic long-duration GRBs have been identified (Bloom et al. 2003; Soderberg et al. 2004, 2006). They too are associated with SNe Ib/c, but their relativistic energy release is a factor of 100 less than that of typical cosmological GRBs and their outflows are significantly less collimated (quasi-spherical). Since they can only be detected at low redshifts where the comparative volume for discovery is low, they are small in total number. But their volumetric rate is inferred to be 10–100 times larger than that of the more distant long-duration GRBs (Soderberg et al. 2006; Cobb et al. 2006; Liang et al. 2007).

More recently, evidence has been growing for a class of GRBs whose total relativistic energy release is at least an order of magnitude above the canonical value of  $10^{51}$  erg (e.g., Cenko et al. 2010 and references therein). Unlike subluminal events, the total energy budget of these hyper-energetic events poses a significant challenge for some progenitor models. In particular, models in which the GRB is powered by a magnetar or a neutrino-driven collapsar are strongly disfavored. On the other hand, collapsars driven by magneto-hydrodynamical (MHD) processes, such as the Blandford-Znajek mechanism (Blandford & Znajek 1977), can naturally accommodate energy budgets as large as  $10^{53}$  erg.

Unfortunately, it has been rather difficult to constrain the beaming-corrected energetics for the hundreds of GRBs detected by the *Swift* satellite (Gehrels et al. 2004). The reasons for this difficulty are now largely understood. First, the relatively narrow energy bandpass (15–150 keV) can miss entirely the peak of the gamma-ray spectrum, making estimates of  $E_{\gamma,\text{iso}}$  highly uncertain. Second, there has been a dearth of measurements of jet opening angles (e.g., Panaitescu 2007; Kocevski & Butler 2008; Liang et al. 2008; Racusin et al. 2009) and well-sampled multi-wavelength GRB afterglows (used to derive the afterglow kinetic energy  $E_{\text{KE}}$ ). *Swift* GRBs are on average more than twice as distant (Jakobsson et al. 2006) and




FIG. 1.— Prompt isotropic gamma-ray energy release ( $E_{\gamma,\text{iso}}$ ) of GRBs. With its soft, narrow bandpass (15–150 keV), *Swift* typically selects events with smaller isotropic energy release but larger opening angles than previous missions, which triggered predominantly in the MeV bandpass (Perna et al. 2003). GRBs detected at GeV energies with the *Fermi*-LAT all fall at the brightest end of the isotropic energy distribution, and must therefore be highly collimated to achieve a canonical beaming-corrected energy release of  $\sim 10^{51}$  erg. References: pre-*Swift*: Amati (2006); *Swift*: Butler et al. (2007); *Fermi*-LAT: Greiner et al. (2009), this work.

therefore significantly fainter ( $\sim 1.5$  mag in the optical; Berger et al. 2005; Kann et al. 2007) than GRBs in previous samples. In large part this is due to selection effects: a combination of bandpass and sensitivity from *Swift* has preferentially selected the faint end of the luminosity function — GRBs with low isotropic energy release but large opening angles (Perna et al. 2003).

With its nearly seven decades in energy coverage (10 keV – 100 GeV), *Fermi* can provide unparalleled constraints on this subsample of the most luminous events. In light of the empirical relation between the peak energy of the gamma-ray spectrum and the isotropic gamma-ray energy release (the  $E_p$ – $E_{\gamma,\text{iso}}$ , or “Amati” relation; Amati 2006), MeV/GeV events detected by either the Gamma-Ray Burst Monitor (GBM, 8 keV – 40 MeV; Meegan et al. 2009) or the Large Area Telescope (LAT, 20 MeV – 300 GeV; Atwood et al. 2009) onboard *Fermi* preferentially select a sample of GRBs with large isotropic energy release (Figure 1). High- $E_{\gamma,\text{iso}}$  events also have brighter X-ray and optical afterglows on average (Nysewander et al. 2009). Follow-up afterglow observations can then determine whether these GRBs are highly beamed events ( $\theta \lesssim 2^\circ$ ) with a typical energy release or true hyper-energetic GRBs.

The *Fermi*-LAT offers a further advantage over previous GRB missions sensitive only at MeV and keV energies by providing strict constraints on the initial Lorentz factor of the relativistic outflow. To avoid  $e^+ - e^-$  pair production (and the accompanying thermal spectrum), the GRB jet must be moving towards the observer with ultra-relativistic speeds (the “compactness” problem; Cavallo & Rees 1978). The higher the energy of the most energetic photon detected from a GRB, the more strict the lower limit on the outflow Lorentz factor will be. Combining the Lorentz factor limits for the most relativistic GRBs with inferred jet opening angles from broadband afterglow models can provide critical diagnostics of the jet acceleration mechanism.

Here we report on broadband (radio, optical, and




FIG. 2.— The broadband radio (blue), optical (red), and X-ray (black) light curve of GRB 090323. The best-fit model is plotted in solid lines (see Table 4 for parameters). The identical model parameters for an isotropic explosion are plotted as the dashed lines. The strength of the possible modulation of the radio afterglow caused by interstellar scintillation (e.g., Frail et al. 2000a) is indicated by the light-blue shaded region. The model provides a reasonable fit in all bandpasses. It is clear that any jet break must occur at  $t > 10$  days, although the upper bound on the jet break time is only weakly constrained.

X-ray) observations of four long-duration GRBs detected by the *Fermi*-LAT at GeV energies: GRBs 090323, 090328, 090902B, and 090926A. For each event we construct afterglow models to constrain the collimation and beaming-corrected energetics, and we compare these LAT events with previous GRBs detected at other energies (i.e., keV energies from *Swift*, and MeV energies from pre-*Swift* satellites). For three of these GRBs, we also present the optical spectra used to determine the afterglow redshift. A more thorough analysis of the host-galaxy properties of these events will be presented in a forthcoming work.

Throughout this paper, we adopt a standard  $\Lambda$ CDM cosmology with  $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_m = 0.27$ , and  $\Omega_\Lambda = 1 - \Omega_m = 0.73$  (Spergel et al. 2007). We define the flux-density power-law temporal and spectral decay indices  $\alpha$  and  $\beta$  as  $f_\nu \propto t^{-\alpha} \nu^{-\beta}$  (e.g., Sari et al. 1998). All quoted uncertainties are  $1\sigma$  (68%) confidence intervals unless otherwise noted.

## 2. OBSERVATIONS

### 2.1. GRB 090323

#### 2.1.1. High-Energy Properties

GRB 090323 was detected by the *Fermi* GBM at 00:02:42.63 on 23 March 2009 (Ohno et al. 2009; UT dates are used throughout this work). In the 8 keV to 40 MeV bandpass of the GBM, the light curve was multip peaked with a duration<sup>16</sup> of  $t_{90} \approx 150$  s. GRB 090323 was also detected at MeV energies by several of the satellites comprising the Inter-Planetary Network (IPN; Hurley et al. 1999), including the Konus instrument on

the *Wind* satellite (Hurley et al. 2009; Golenetskii et al. 2009c).

Examining only the first 70 s of GBM data<sup>17</sup>, van der Horst & Xin (2009) find that the prompt spectrum is well described by a power law having an exponential cutoff at high energies with  $\alpha = -0.89 \pm 0.03$  and  $E_p = 697 \pm 51$  keV. The resulting fluence in the 8–10<sup>3</sup> keV (observer frame) bandpass is  $f_\gamma = (1.00 \pm 0.01) \times 10^{-4} \text{ erg cm}^{-2}$ . The Konus-*Wind* instrument was able to measure the high-energy spectrum over the entire duration of the MeV emission. Fitting a Band model (Band et al. 1993) to their spectrum, Golenetskii et al. (2009c) find a reasonable fit with  $\alpha = -0.96^{+0.12}_{-0.09}$ ,  $\beta = -2.09^{+0.16}_{-0.22}$ ,  $E_p = 416^{+76}_{-73}$  keV, and  $f_\gamma = 2.02^{+0.28}_{-0.25} \times 10^{-4} \text{ erg cm}^{-2}$  (20–10<sup>4</sup> keV observer-frame bandpass).

In order to facilitate direct comparisons between events, we must transform this high-energy fluence to a common (rest-frame) bandpass (i.e., a “k”-correction; Bloom et al. 2001). Here we adopt the rest-frame 1–10<sup>4</sup> keV bandpass, as this encompasses the full range of peak energies observed from GRBs (Band et al. 1993) without (for most pre-*Fermi* satellites) requiring large extrapolations outside the observed bandpass. At  $z = 3.568$  (§ 2.1.3), The Konus-*Wind* measurement corresponds to a prompt fluence of  $f_\gamma = (1.50 \pm 0.20) \times 10^{-4} \text{ erg cm}^{-2}$  in the 1–10<sup>4</sup> keV rest-frame bandpass. Given the limited temporal coverage of the GBM spectrum, we shall adopt this value for the prompt fluence of GRB 090323 for the remainder of this work.

In addition to the detection at MeV energies by the GBM, GRB 090323 was also detected at GeV energies

<sup>16</sup> It is customary to report GRB durations measured as the time between the arrival of 5% and 95% of the background-subtracted fluence. This quantity is referred to as  $t_{90}$ .

<sup>17</sup> After this point, the *Fermi* spacecraft slewed to position the GRB at the center of the LAT field of view, causing rapid changes in the GBM background level.




FIG. 3.— GMOS-S optical spectrum of the afterglow of GRB 090323. The broad absorption feature at  $\lambda \approx 5570 \text{ \AA}$  results from  $\text{Ly}\alpha$  in the GRB host galaxy. We identify a number of strong, narrow absorption features redward of  $\text{Ly}\alpha$  from the GRB host galaxy at a common redshift of  $3.568 \pm 0.004$  (black annotations). The emission blueward of  $\text{Ly}\alpha$  is strongly affected by the  $\text{Ly}\alpha$  forest. We identify two additional intervening absorbers, based on  $\text{Mg II } \lambda\lambda 2796, 2803$  ( $z = 2.101$ ; blue annotations) and  $\text{C IV } \lambda\lambda 1548, 1550$  ( $z = 3.379$ ; red annotations). Areas of the spectrum affected by telluric absorption in the atmosphere of the Earth are marked with crossed circles. Because of the increased noise in the NIR due to fringing, we have cut off the plot at  $\lambda = 9000 \text{ \AA}$ .

by the *Fermi* LAT. Like several previous GRBs observed at GeV energies (e.g., Hurley et al. 1994; González et al. 2003; Giuliani et al. 2008; Abdo et al. 2009b), the GeV emission began several seconds after the MeV emission is detected, and remains significant long ( $\sim 1000 \text{ s}$ ) after the MeV component has faded beyond detectability (Ohno et al. 2009). The highest energy photon from the direction of GRB 090323 had an energy of  $E \approx 7.5 \text{ GeV}$  (Piron et al. 2009). However, this photon arrived after the prompt MeV emission ended, and so its origin (i.e., prompt emission or afterglow, or, in physical terms, internal or external shock-related) is somewhat uncertain. The highest energy photon detected by the LAT during the formal MeV  $t_{90}$  measurement had an energy of  $E \approx 500 \text{ keV}$  (at  $t \approx 90 \text{ s}$ ; Piron et al. 2009).

### 2.1.2. Afterglow Observations

The *Swift* satellite began observations of the field of GRB 090323 with the onboard X-ray Telescope (XRT; Burrows et al. 2005b) at 19:27 on 23 March 2009 ( $\sim 19.4$  hours after the burst; Kennea et al. 2009a). A candidate afterglow was promptly identified at  $\alpha = 12^{\text{h}}42^{\text{m}}50.^{\text{s}}26$ ,  $\delta = +17^{\circ}03'14''.2$ , with a 90% containment radius of  $2''.7$  (J2000.0). The XRT continued to monitor the evolution of the fading counterpart for the following two weeks. We have obtained the XRT light curve from the online compilation of N.R.B.<sup>18</sup>; the resulting evolution is plotted in Figure 2.

The optical afterglow of GRB 090323 was discovered shortly thereafter by GROND (Udike et al. 2009b). The optical/NIR spectral energy distribution, constructed from simultaneous imaging in the  $g'r'i'z'JHK$

filters, implied a spectral steepening around the observed  $g'$  band. Associating this break with absorption from  $\text{Ly}\alpha$  in the GRB host galaxy, Udike et al. (2009b) derive a photometric redshift of  $4.0 \pm 0.3$  for GRB 090323.

We began observations of the afterglow of GRB 090323 with the automated Palomar 60 inch (1.5 m) telescope (P60; Cenko et al. 2006) beginning on 2009 March 24 (Cenko & Perley 2009). Images were obtained in the Sloan  $r'$  and  $i'$  filters and individual frames were automatically reduced using our custom IRAF<sup>19</sup> software pipeline. To increase the signal-to-noise ratio (S/N), individual frames were astrometrically aligned using the Scamp software package and coadded using Swarp<sup>20</sup>. We used aperture photometry to extract the flux of the afterglow from these coadded frames with the aperture radius roughly matched to the full width at half-maximum intensity (FWHM) of the point-spread function (PSF). Aperture magnitudes were then calibrated relative to field sources from the Sloan Digital Sky Survey Data Release 7 (Abazajian et al. 2009). Imaging continued on subsequent nights with P60 until the afterglow was below our detection threshold. The results of this monitoring campaign, uncorrected for foreground Galactic extinction ( $E[B - V] = 0.025 \text{ mag}$ ; Schlegel et al. 1998), are presented in Table 9.

We obtained additional imaging of the field of GRB 090323 with the two Gemini Multi-Object Spectrographs (GMOS-N and GMOS-S; Hook et al. 2004). Images were taken at both Gemini North and Gemini South in the Sloan  $r'$  and  $i'$  filters, and they were reduced using

<sup>18</sup> <http://astro.berkeley.edu/~nat/swift/>; see Butler & Kocevski (2007) for details.

<sup>19</sup> IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

<sup>20</sup> See <http://astromatic.iap.fr>.

TABLE 1  
RADIO OBSERVATIONS OF GRB 090323

Date (UT)	$\Delta t$ (days)	$\nu$ (GHz)	$f_\nu$ ( $\mu$ Jy)	Facility
Mar 26.38	3.38	8.46	$27 \pm 38$	VLA
Mar 27.38	4.38	8.46	$225 \pm 35$	VLA
Mar 27.99	4.99	4.9	$105 \pm 24$	WSRT <sup>a</sup>
Mar 28.43	5.43	8.46	$100 \pm 40$	VLA
Mar 29.16	6.16	8.46	$157 \pm 31$	VLA
Mar 30.18	7.18	8.46	$219 \pm 39$	VLA
Mar 31.32	8.32	8.46	$281 \pm 38$	VLA
Apr 1.30	9.30	8.46	$164 \pm 35$	VLA
Apr 3.29	11.29	8.46	$166 \pm 27$	VLA
Apr 3.28	11.28	4.86	$110 \pm 45$	VLA
Apr 4.41	12.41	8.46	$183 \pm 35$	VLA
Apr 5.14	13.14	8.46	$123 \pm 29$	VLA
Apr 6.28	14.28	8.46	$312 \pm 27$	VLA
Apr 7.42	15.42	8.46	$43 \pm 27$	VLA
Apr 9.44	17.44	8.46	$127 \pm 30$	VLA
Apr 10.42	18.42	8.46	$295 \pm 27$	VLA
Apr 14.09	22.09	8.46	$178 \pm 56$	VLA
Apr 17.04	25.04	8.46	$167 \pm 33$	VLA
Apr 25.46	33.46	8.46	$78 \pm 32$	VLA
May 3.41	41.41	8.46	$77 \pm 31$	VLA
May 11.35	49.35	8.46	$61 \pm 29$	VLA
Aug 1.18	131.18	8.46	$-13 \pm 27$	VLA

<sup>a</sup> Reference: van der Horst (2009).

the IRAF *gemin*i package. Photometry was performed with the same methodology used for the P60 imaging of the field, and the resulting measurements are shown in Table 9.

Additionally, we have compiled optical and NIR measurements of the afterglow of GRB 090323 from the GCN<sup>21</sup> Circulars and included these in Table 9. The majority of the late-time measurements (of most interest to us for modeling purposes) were obtained in the *R* band. We note that most of these measurements were calibrated with respect to a single object (1070-0238439) from the USNO-B1 catalog (Monet et al. 2003), following Kann et al. (2009b). Using the SDSS observations of this field and the filter transformations of Jordi et al. (2006), we find an *R*-band magnitude for this source of  $R = 17.31$  mag (compared to an assumed value of 17.15 mag from Kann et al. 2009b). We therefore offset all reported *R*-band magnitudes from the GCN Circulars by +0.16 mag.

Finally, we observed the afterglow of GRB 090323 with the Very Large Array<sup>22</sup> (VLA) beginning a few days after the initial burst trigger. The flux-density scale was tied to 3C 286 or 3C 147 and the phase was measured by switching between the GRB and a nearby bright point-source calibrator. To maximize sensitivity, the full VLA continuum bandwidth (100 MHz) was recorded in two 50 MHz bands. Data reduction was carried out following standard practice in the *AIPS* software package.

Our initial VLA detection of GRB 090323 was reported by Harrison et al. (2009). GRB 090323 was also detected at radio wavelengths by the Westerbork Synthesis Radio Telescope (WSRT; van der Horst 2009). The full set of VLA measurements is listed in Table 1. In order to improve the S/N and to reduce the modulation of the light

curve caused by interstellar scintillation (e.g., Frail et al. 2000a), we binned the data from adjacent epochs. These binned points were used for our afterglow modeling (§ 3) and are plotted in Figure 2.

### 2.1.3. Optical Spectroscopy

We began spectroscopic observations of the optical afterglow of GRB 090323 with GMOS-S on 24 March 2009 ( $\sim 29.93$  hours after the GBM trigger; Chornock et al. 2009). We first obtained  $2 \times 600$  s spectra with the B600 grating and a central wavelength of 6000 Å, providing coverage over the range  $\sim 4500$ –7500 Å with a resolution of 2.5 Å. Immediately following these exposures, we obtained  $2 \times 600$  s spectra with the R400 grating and a central wavelength of 8000 Å, providing coverage of  $\sim 6000$ –10000 Å with a resolution of 5.0 Å.

The spectra were reduced in a standard manner using routines from the IRAF *gemin*i and *specred* packages (see, e.g., Cenko et al. 2008 and references therein for details). Wavelength calibration was performed relative to CuAr lamps and then adjusted based on measured night-sky emission lines. The resulting root-mean square wavelength uncertainty was  $\lesssim 0.25$  Å for the spectra with the B600 grating and  $\lesssim 0.40$  Å for the spectra taken with the R400 grating. Telluric features were removed using the continuum from well-exposed spectrophotometric standard stars (e.g., Wade & Horne 1988; Matheson et al. 2000).

Flux calibration was performed relative to the spectrophotometric standard star LTT 7379 (Stone & Baldwin 1983; Baldwin & Stone 1984). We caution, however, that the standard-star observations were conducted on different nights from the GRB observations (31 August 2009 for the B600 grating; 4 August 2009 for the R400 grating), so the absolute flux calibration is somewhat uncertain (estimated to be  $\sim 30\%$ ).

The resulting spectrum of the afterglow of GRB 090323 is shown in Figure 3. The broad absorption feature at  $\lambda \approx 5570$  Å is produced by Ly $\alpha$  at  $z \approx 3.6$ . The spectrum blueward of this wavelength is dominated by absorption from the Ly $\alpha$  forest, while the lack of Ly $\alpha$  forest features redward of this transition indicates that it corresponds to the redshift of the GRB host galaxy. Unfortunately, the GMOS CCD chip gap over the range 5500–5525 Å precludes an accurate measurement of the Ly $\alpha$  profile and hence a determination of the H I column density.

Redward of Ly $\alpha$ , we find a series of strong (rest-frame equivalent width  $W_0 \geq 1$  Å) absorption features superposed on a relatively flat, featureless continuum. In particular, we identify Si II  $\lambda 1260$ , Si II\*  $\lambda 1264$ , O I  $\lambda 1302$ , Si II  $\lambda 1304$ , Si II\*  $\lambda 1309$ , C II  $\lambda 1334$ , C II\*  $\lambda 1335$ , Si IV  $\lambda \lambda 1393, 1402$ , Si II  $\lambda 1526$ , Si II\*  $\lambda 1533$ , and C IV  $\lambda \lambda 1548, 1550$  at  $z = 3.568 \pm 0.004$ . Excited fine structure lines are indicative of UV pumping and further suggest these features arise in the host galaxy of GRB 090323. The strongest features exhibit a complex velocity structure that is only marginally resolved at the resolution of our B600 spectra. We also detect intervening absorption systems of C IV  $\lambda \lambda 1548, 1550$  at  $z = 3.379$  and Mg II  $\lambda \lambda 2796, 2803$  at  $z = 2.101$ . A more thorough analysis of the host-galaxy properties of GRB 090323 will be presented in a subsequent work.

<sup>21</sup> See [http://gcn.gsfc.nasa.gov/gcn3\\_archive.html](http://gcn.gsfc.nasa.gov/gcn3_archive.html).

<sup>22</sup> The Very Large Array is operated by the National Radio Astronomy Observatory, a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.






FIG. 4.— The broadband radio (blue), UV (red), and X-ray (black) light curve of GRB 090328. The best-fit model is plotted in solid lines (see Table 5 for parameters). The identical model parameters for an isotropic explosion are plotted as the dashed lines. The radio light curve is not very well fit at early times, although it likely suffers from strong interstellar scintillation (light-blue shaded region). In this case a jet break is required by the data to fall at  $t \gtrsim 10$  days.

At  $z = 3.568$ , the isotropic prompt energy release from GRB 090323 in the rest-frame  $1\text{--}10^4$  keV bandpass is  $E_{\gamma, \text{iso}} = (3.99 \pm 0.53) \times 10^{54}$  erg. Using the formulation of Lithwick & Sari (2001), the lower limit on the outflow Lorentz factor (assuming a nonthermal spectrum up to  $E_{\text{obs}} \approx 500$  keV) is  $\Gamma_0 \gtrsim 600$ .

## 2.2. GRB 090328

### 2.2.1. High-Energy Properties

GRB 090328 triggered the *Fermi* GBM at 09:36:46 on 28 March 2009 (McEnery et al. 2009). In the GBM bandpass, the light curve is multi-peaked with a duration of  $t_{90} \approx 70$  s. GRB 090328 was also detected at MeV energies by several IPN satellites (Golenetskii et al. 2009a), including *Konus-Wind* (Golenetskii et al. 2009d).

A spectrum consisting of the first  $\sim 30$  s of GBM data (containing the brightest part of the high-energy light curve) is well fit by a Band function with  $\alpha = -0.93 \pm 0.02$ ,  $\beta = -2.2 \pm 0.1$ ,  $E_p = 653 \pm 45$  keV (Rau et al. 2009). The resulting fluence in the 8 keV to 40 MeV (observer frame) bandpass is  $f_\gamma = (9.5 \pm 1.0) \times 10^{-5}$  erg cm $^{-2}$ . These results are all consistent with the analogous values derived by the *Konus-Wind* instrument, which include data out to  $t_0 + 73$  s. Given the wider bandpass of the GBM and the higher precision of its fluence measurement, we adopt this value for remainder of this work. Using the observed redshift of 0.736 (§ 2.1.3), we find a prompt fluence of  $f_\gamma = (7.3 \pm 0.8) \times 10^{-5}$  erg cm $^{-2}$  in the  $1\text{--}10^4$  keV rest-frame bandpass.

GRB 090328 was also detected at GeV energies by the *Fermi*-LAT. Much like GRB 090323, flux in the LAT band from GRB 090328 is detected out to  $\sim 900$  s after the GBM trigger (Cutini et al. 2009). Many photons with energies above 1 GeV are detected from the direction of GRB 090328; however, they are all detected well after the end of the MeV emission (Cutini et al.

2009; Piron et al. 2009). The highest energy photon detected from the direction of GRB 090328 was measured at  $\sim 5$  GeV ( $t_0 + 798$  s), while the most energetic photon detected during the prompt emission had  $E \approx 700$  keV at  $t_0 + 60$  s (Piron et al. 2009).

### 2.2.2. Afterglow Observations

The *Swift* XRT began target-of-opportunity observations of GRB 090328 at 01:26 on 29 March 2009 ( $\sim 15.9$  hours after the GBM trigger). A fading X-ray source at  $\alpha = 6^{\text{h}}02^{\text{m}}39.^{\text{s}}58$ ,  $\delta = -41^\circ 52' 57''.5$  (J2000.0;  $2''.0$  containment radius) was promptly identified as the X-ray afterglow (Kennea 2009; Kennea et al. 2009b). The resulting X-ray light curve, showing the afterglow evolution over the following 10 days, is plotted in Figure 4.

The Ultraviolet Optical Telescope (UVOT; Roming et al. 2005) onboard *Swift* began settled observations of the field of GRB 090328 approximately 16 hours after the *Fermi* trigger. The optical afterglow, detected in both the *U*-band and white filters, was identified shortly thereafter (Oates 2009). UVOT observations of the afterglow continued for  $\sim 2$  weeks, almost exclusively in these two filters. We have downloaded the UVOT *U*-band data from the HEASARC archive<sup>23</sup> and conducted photometry with the images following the technique described by Li et al. (2006). Given the large uncertainty associated with flux calibration for the white filter, we have not included data taken in this band. The results of this analysis, not including a correction for foreground Galactic extinction ( $E[B - V] = 0.057$  mag; Schlegel et al. 1998), are shown in Table 10.

We obtained a single  $i'$  image of the afterglow of GRB 090328 with GMOS-S on 29 April 2009. This image was reduced in an identical manner to our observa-

<sup>23</sup> See <http://heasarc.gsfc.nasa.gov>.




FIG. 5.— GMOS-S optical spectrum of the afterglow of GRB 090328. We identify a series of forbidden O emission lines and Fe, Mg, and Ca absorption features at a common redshift of  $0.7357 \pm 0.0004$ . We have only plotted the wavelength range 4500–9000 Å, due to the lower S/N in the blue (spectrograph throughput) and the red (fringing).

TABLE 2  
VLA RADIO OBSERVATIONS OF  
GRB 090328

Date (UT)	$\Delta t$ (days)	$\nu$ (GHz)	$f_\nu$ ( $\mu$ Jy)
Mar 30.02	1.62	8.46	$172 \pm 57$
Mar 30.99	2.59	8.46	$337 \pm 60$
Apr 1.02	3.62	8.46	$783 \pm 57$
Apr 3.03	5.63	8.46	$195 \pm 34$
Apr 4.01	6.61	8.46	$674 \pm 155$
Apr 5.09	7.69	8.46	$214 \pm 41$
Apr 6.02	8.62	8.46	$523 \pm 57$
Apr 6.97	9.57	8.46	$809 \pm 39$
Apr 11.14	13.74	8.46	$603 \pm 67$
Apr 14.00	16.60	8.46	$643 \pm 59$
Apr 16.99	19.59	8.46	$886 \pm 58$
Apr 24.97	27.57	8.46	$410 \pm 59$
May 9.96	42.56	8.46	$78 \pm 43$
May 11.86	44.46	8.46	$21 \pm 64$
May 17.92	50.52	8.46	$-20 \pm 40$

tions of GRB 090323, and the zero point was calculated using predetermined values from the Gemini website<sup>24</sup>. We have also included the simultaneous GROND optical and NIR measurements from Updike et al. (2009a) in our modeling, and therefore display them in Table 10 as well.

Finally, we began observing the afterglow of GRB 090328 with the VLA on 30 March 2009 (Frail et al. 2009) and continued for nearly two months. The data were reduced in a manner identical to that described in § 2.1.2. The results of our radio campaign are displayed in Table 2 (individual epochs) and plotted in Figure 4 (combined epochs).

### 2.2.3. Optical Spectroscopy

We began spectroscopic observations of GRB 090328 with GMOS-S at 00:05 on 30 April 2009 ( $\sim 38.5$  hours after the GBM trigger; Cenko et al. 2009). We obtained  $2 \times 1500$  s spectra with the R400 grating, the first with a central wavelength of 6000 Å (providing coverage of  $\sim$

4000–8000 Å), and the second with a central wavelength of 8000 Å (providing coverage of  $\sim 6000$ –10000 Å). The spectra were reduced in the manner described in § 2.1.3. Flux calibration was performed relative to the standard star LTT 7379 taken with identical instrumental setups on 4 August 2009.

The resulting spectrum of GRB 090328 is shown in Figure 5. Superposed on a relatively flat continuum, we identify strong emission lines of oxygen ([O II]  $\lambda 3727$ , [O III]  $\lambda\lambda 4959, 5007$ ) as well as a series of narrow absorption features (Fe II  $\lambda 2586$ , Fe II  $\lambda 2600$ , Mg II  $\lambda\lambda 2796, 2803$ , Mg I  $\lambda 2852$ , and Ca II  $\lambda\lambda 3934, 3969$ ), all at  $z = 0.7357 \pm 0.0004$ .

At this redshift, the prompt isotropic gamma-ray energy release from GRB 090328 in the 1–10<sup>4</sup> keV bandpass is  $E_{\gamma, \text{iso}} = (1.34 \pm 0.14) \times 10^{53}$  erg. Using the formulation of Lithwick & Sari (2001), the lower limit on the outflow Lorentz factor (assuming a nonthermal spectrum up to  $E_{\text{obs}} \approx 700$  keV) is  $\Gamma_0 \gtrsim 200$ .

## 2.3. GRB 090902B

### 2.3.1. High-Energy Properties

At 11:05:08.31 on 2 September 2009, the *Fermi*-GBM triggered and located GRB 090902B (Bissaldi & Connaughton 2009). In the GBM bandpass, the light curve consisted of a bright, multi-peaked pulse with a duration  $t_{90} \approx 21$  s. GRB 090902B was also detected in a similar bandpass by *Suzaku*-WAM (Terada et al. 2009).

Furthermore, GRB 090902B was bright enough to be detected at GeV energies by the *Fermi*-LAT (de Palma et al. 2009). Like the other events in our sample, LAT emission was seen out to  $t_0 + 1000$  s, at late times decaying like a power law with temporal index  $\alpha_{\text{LAT}} \approx 1.5$  (Abdo et al. 2009a). GRB 090902B is responsible for the highest energy photon detected from a GRB to date, with  $E = 33.4^{+3.7}_{-3.5}$  GeV at  $t_0 + 82$  s. Within the MeV prompt emission phase, the highest energy photon was measured at  $E = 11.16^{+1.48}_{-0.58}$  GeV (Abdo et al. 2009a).

Unlike the other events considered here, the prompt high-energy spectrum of GRB 090902B is not adequately

<sup>24</sup> <http://www.gemini.edu/sciops/instruments/gmos/?q=sciops/instruments/gmos>.




FIG. 6.— The broadband radio (blue), optical (red), UV (cyan), and X-ray (black) light curves of GRB 090902B. The best-fit model is plotted in solid lines (see Table 6 for parameters). The identical model parameters for an isotropic explosion are plotted as the dashed lines. The strength of the possible modulation of the radio afterglow caused by interstellar scintillation (e.g., Frail et al. 2000a) is indicated by the light-blue shaded region. As suggested by Pandey et al. (2010), the early ( $t \lesssim 0.3$  day) optical (and, to a lesser extent, radio) data are significantly larger than predicted by our forward-shock afterglow models. The most likely explanation is the presence of reverse-shock emission. The lack of a bright radio afterglow at late times suggests the presence of a jet break at  $t \approx 6$  days.

described by a single Band function. While the 100–1000 keV bandpass is reasonably well fit by such a model, the spectrum exhibits excess emission at both low ( $\lesssim 100$  keV) and high ( $\gtrsim 10$  MeV) energies. Abdo et al. (2009a) have suggested that the high-energy spectrum of GRB 090902B can be reproduced as the sum of two components: a Band function peaking at  $\sim 700$  keV, and a single power-law (photon index  $\Gamma \equiv \beta + 1 = 1.93$ ) extending over the entire GBM+LAT bandpass. In this model, the power-law component accounts for  $\sim 24\%$  of the total 10 keV to 10 GeV fluence. The physical mechanism responsible for this complex spectrum is still not entirely understood; possible explanations include a hadronic origin [either proton synchrotron radiation (Razzaque et al. 2009) or photohadronic interactions (Asano et al. 2009)] or thermal emission from the jet photosphere (Mészáros & Rees 2000; Ryde 2004; Ryde et al. 2010).

Using their two-component spectrum, Abdo et al. (2009a) report a total fluence of  $f_\gamma = (4.59 \pm 0.05) \times 10^{-4}$  erg cm $^{-2}$  in the 8 keV to 30 GeV (observer frame) bandpass. At a redshift of 1.8229 (§ 2.3.3), this corresponds to a prompt fluence of  $(3.83 \pm 0.05) \times 10^{-4}$  erg cm $^{-2}$  in the 1–10 $^4$  keV rest-frame bandpass.

### 2.3.2. Afterglow Observations

The *Swift* XRT began target-of-opportunity observations of the field of GRB 090902B at 23:36 on 2 September 2009 ( $\sim 12.5$  hr after the GBM trigger). A fading X-ray source at  $\alpha = 17^{\text{h}}39^{\text{m}}45.^{\text{s}}26$ ,  $\delta = +27^{\circ}19'28''.1$  (J2000.0;  $2''.1$  containment radius) was promptly identified as the X-ray afterglow (Kennea & Stratta 2009; Evans 2009; Stratta et al. 2009). The resulting X-ray light curve is shown in Figure 6.

The *Swift* UVOT began concurrently observing the field of GRB 090902B and first reported the detection of a candidate optical afterglow consistent with the X-ray position (Swenson & Siegel 2009). Subsequent observations revealed that the candidate had faded, confirming its association with GRB 090902B (Swenson & Stratta 2009). In Table 11 we present UVOT *U*-band observations of GRB 090902B, reduced in an identical manner to those described in § 2.2.2.

We obtained a single  $r'$  image of the afterglow of GRB 090902B with GMOS-N on 3 September 2009. This image was reduced in the same manner as those of the other events, and the resulting photometry is presented in Table 11. We have included optical and NIR photometry of GRB 090902B from Pandey et al. (2010) in our modeling, and therefore display them in Table 11 as well.

Finally, we began observing the afterglow of GRB 090902B with the VLA on 3 September 2009 (Chandra & Frail 2009) and continued for  $\sim 5$  months. The data were reduced as described in § 2.2.2. The results of our radio campaign are displayed in Tables 3 (individual epochs) and plotted in Figure 6 (combined epochs). We have also included the early radio detection at 4.8 GHz reported by the WSRT (van der Horst et al. 2009) in Table 3.

### 2.3.3. Optical Spectroscopy

We began spectroscopic observations of GRB 090902B with GMOS-N at 06:29 on 3 September 2009 ( $\sim 19.4$  hours after the GBM trigger; Cucchiara et al. 2009). We obtained  $2 \times 900$  s exposures, both with the R400 grating and a central wavelength of 6000 Å, providing coverage of  $\sim 4000$ –8000 Å. The spectra were reduced as described in § 2.1.3 and § 2.2.3. Flux calibration was per-






FIG. 7.— GMOS-N optical spectrum of the afterglow of GRB 090902B. We identify a series of absorption features from neutral and singly ionized Mg, Mn, Fe, and Cr at a common redshift of  $1.8229 \pm 0.0004$ . We have not plotted the shortest wavelengths due to the decreased spectrograph throughput.

TABLE 3  
RADIO OBSERVATIONS OF GRB 090902B

Date (UT)	$\Delta t$ (days)	$\nu$ (GHz)	$f_\nu$ ( $\mu\text{Jy}$ )	Facility
2009 Sep 3.77	1.31	4.8	$111 \pm 28$	WSRT <sup>a</sup>
2009 Sep 3.94	1.48	8.46	$141 \pm 39$	VLA
2009 Sep 7.05	4.59	8.46	$13 \pm 31$	VLA
2009 Sep 8.05	5.59	8.46	$130 \pm 34$	VLA
2009 Sep 10.15	7.69	8.46	$10 \pm 32$	VLA
2009 Sep 11.05	8.59	8.46	$80 \pm 32$	VLA
2009 Sep 13.10	10.64	8.46	$99 \pm 31$	VLA
2009 Sep 14.14	11.60	8.46	$71 \pm 33$	VLA
2009 Sep 18.04	15.50	8.46	$52 \pm 32$	VLA
2009 Sep 19.00	16.46	8.46	$89 \pm 36$	VLA
2009 Sep 25.09	22.51	8.46	$26 \pm 29$	VLA
2009 Sep 27.08	24.50	8.46	$67 \pm 29$	VLA
2009 Oct 7.01	34.43	8.46	$38 \pm 28$	VLA
2009 Oct 9.01	36.43	8.46	$66 \pm 27$	VLA
2009 Oct 11.81	39.23	8.46	$21 \pm 31$	VLA
2009 Nov 6.90	65.44	8.46	$9 \pm 20$	VLA
2009 Nov 7.77	66.31	8.46	$22 \pm 19$	VLA
2009 Nov 9.00	67.54	8.46	$48 \pm 19$	VLA
2009 Nov 14.88	73.42	8.46	$31 \pm 21$	VLA
2010 Mar 20.62	199.16	8.46	$18 \pm 16$	VLA

<sup>a</sup> Reference: van der Horst et al. (2009).

formed relative to spectra of the standard star Feige 34 (Oke 1990) taken with the same instrumental setup as on 1 May 2008.

The resulting spectrum of GRB 090902B is shown in Figure 7. Super-imposed on a smooth power-law continuum, we identify strong absorption features corresponding to Mg I  $\lambda 2852$ , Mg II  $\lambda\lambda 2796, 2803$ , Mn II  $\lambda 2606$ , Fe II  $\lambda 2600$ , Mn II  $\lambda 2594$ , Fe II  $\lambda 2586$ , Mn II  $\lambda 2576$ , Fe I  $\lambda 2484$ , Fe II  $\lambda 2382$ , Fe II  $\lambda 2344$ , Fe II  $\lambda 2260$ , Fe I  $\lambda 2167$ , Cr II  $\lambda\lambda 2056, 2062$ , and Mg I  $\lambda 2026$ , all at a common redshift of  $1.8229 \pm 0.0004$ .

The lack of evidence for Ly $\alpha$  absorption down to  $\lambda \lesssim 4000 \text{ \AA}$  implies an upper bound on the redshift of the GRB host galaxy of  $z \lesssim 2.3$ . Together with the strength of the above features (in particular Fe I  $\lambda 2167$ ), we consider it quite likely that the observed system at

$z = 1.8229$  derives from the GRB host galaxy.

At this redshift, the prompt isotropic gamma-ray energy release in the  $1\text{--}10^4 \text{ keV}$  rest-frame bandpass is  $E_{\gamma, \text{iso}} = (3.20 \pm 0.04) \times 10^{54} \text{ erg}$ . Assuming a nonthermal spectrum up to  $E_{\text{obs}} = 11 \text{ GeV}$ , Abdo et al. (2009a) have determined a lower limit to the initial outflow Lorentz factor of  $\Gamma_0 \gtrsim 1000$ .

## 2.4. GRB 090926A

### 2.4.1. High-Energy Properties

GRB 090926A triggered the GBM on *Fermi* at 04:20:26.99 on 26 September 2009 (Bissaldi 2009). The light curve consists of a single pulse with duration  $t_{90} \approx 20 \text{ s}$  in the GBM bandpass. The prompt emission was sufficiently bright to trigger several additional satellite instruments, including *Suzaku*-WAM (Noda et al. 2009), *Konus-Wind* (Golenetskii et al. 2009b), and RT-2 on CORONAS-PHOTON (Chakrabarti et al. 2009).

Simultaneously fitting both the GBM and LAT (see below) data from  $t_0$  to  $t_0 + 21 \text{ s}$ , Bissaldi et al. (2009) report that the spectrum is reasonably well modeled by a Band function with  $\alpha = -0.693 \pm 0.009$ ,  $\beta = -2.342 \pm 0.011$ , and  $E_p = 268 \pm 4 \text{ keV}$ . The corresponding 10 keV to 10 GeV fluence (observer frame) is  $f_\gamma = (2.47 \pm 0.03) \times 10^{-4} \text{ erg cm}^{-2}$ . These results differ only slightly from the values reported from other satellites. At  $z = 2.1062$  (Malesani et al. 2009), the corresponding prompt fluence in the rest-frame  $1\text{--}10^4 \text{ keV}$  bandpass is  $(1.73 \pm 0.03) \times 10^{-4} \text{ erg cm}^{-2}$ .

GRB 090926A was also detected by the *Fermi*-LAT. The emission in the LAT bandpass lasted at least 400 s, with possible indications of significant flux out to  $\sim 1000 \text{ s}$  after the GBM trigger. Many photons with  $E > 1 \text{ GeV}$  were detected from the position of GRB 090926A, with the highest energy measured of 20 GeV at  $t_0 + 26 \text{ s}$  (Uehara et al. 2009).

### 2.4.2. Afterglow Observations

The *Swift* XRT began observing the field of GRB 090926A at 17:17 on 26 September 2009 ( $\sim$




FIG. 8.— Early optical and X-ray afterglow of GRB 090926A. After an initial decline in the  $U$  and  $V$  filters, the afterglow experiences a prominent rebrightening in all filters, peaking at  $t \approx 1$  day. Such behavior is difficult to reconcile with standard afterglow theory, and may suggest a late-time ( $t \gg \Delta t_{\text{GRB}}$ ) injection of energy from the central engine (Rees & Meszaros 1998).

13 hours after the GBM trigger). A fading X-ray counterpart at  $\alpha = 23^{\text{h}}33^{\text{m}}36.^{\text{s}}18$ ,  $\delta = -66^{\circ}19'25''.9$  (J2000.0,  $1''.5$  containment radius) was promptly identified in the XRT data (Vetere et al. 2009; Vetere 2009). The XRT observed GRB 090926A for the next 3 weeks, and we plot the X-ray light curve in Figures 8 and 9.

Under the control of Skynet, four of the 16 inch diameter PROMPT telescopes (Reichart et al. 2005) at Cerro Telolo Inter-American Observatory (CTIO) observed the *Fermi*-LAT localization of GRB 090926A beginning 19.0 hours after the GBM trigger in the  $B$ -,  $V$ -,  $R$ -, and  $I$ -band filters. Within the *Swift*-XRT localization, we identified an uncatalogued and fading source as the optical afterglow of GRB 090926A (Haislip et al. 2009b,a). Individual images were automatically reduced using our custom, IRAF-based reduction pipeline and then astrometrically aligned and stacked. We subsequently measured the afterglow flux with aperture photometry, where the inclusion radius was approximately matched to the FWHM of the PSF. PROMPT continued to observe the field for ten more nights (Haislip et al. 2009e,c,d).

We also obtained 3 epochs of  $I$ - and  $J$ -band imaging of the afterglow of GRB 090926A using the ANDICAM (A Novel Dual Imaging CAMera) instrument mounted on the 1.3 m telescope at CTIO<sup>25</sup>. This telescope is operated as part of the Small and Moderate Aperture Research Telescope System (SMARTS) consortium<sup>26</sup>. Each epoch consisted of 6 individual 360 s  $I$ -band observations and 30 individual 60 s  $J$ -band observations. Between optical exposures, the telescope was slightly offset and the individual  $J$ -band exposures were additionally dithered via an internal tilting mirror system. Standard data reduction was performed on these images, including cosmic ray rejection, overscan bias subtraction, zero subtraction, flat fielding and sky subtraction to correct for the NIR




FIG. 9.— Late-time afterglow and model of GRB 090926A. Both the X-ray and optical ( $g'r'i'$ ) bandpasses exhibit a steepening decline at  $t \approx 10$  days, strongly indicative of a jet break.

background and the  $I$ -band fringing. For each epoch, the individual images were then aligned and averaged to produce a single frame in each band with summed exposure times of 36 minutes in  $I$  and 30 minutes in  $J$ .

Relative aperture photometry was performed on the SMARTS data, in comparison with a number of nonvariable sources in the field of GRB 090926A. The  $I$ -band field was photometrically calibrated by comparison (on photometric nights) with Landolt standard stars in the field of T Phe (Landolt 1992).  $J$ -band photometric calibration was performed using 2MASS (Skrutskie et al. 2006) field stars.

SMARTS  $BVRI$  observations of the field of GRB 090926A were also obtained on two photometric nights a few months after the GRB occurred (16 and 18 December 2009). For these observations, total summed exposure times amounted to 180 s in  $BRI$  and 120 s in  $V$ . The absolute photometry of the field was again established based on same-night observations of the T Phe Landolt standard stars. These observations were then used to provide absolute calibration for the PROMPT observations of GRB 090926A.

We obtained additional late-time imaging of the field of GRB 090926A on 19 October 2009 with GMOS-S on Gemini South. A total of 600 s of exposure time was obtained in the Sloan  $g'$ -,  $r'$ -, and  $i'$ -band filters. The data were reduced in the manner described in § 2.1.2, and calibrated in the same way as the PROMPT and SMARTS observations.

The results of our optical and NIR monitoring campaign of the afterglow of GRB 090926A, uncorrected for the modest amount of Galactic extinction [ $E(B - V) = 0.024$  mag; Schlegel et al. 1998], are shown in Table 12 and Figures 8 and 9.

Finally, the field of GRB 090926A was observed in the radio (5.5 GHz) on 1 October 2009 with the Australia Telescope Compact Array (ATCA). No source was detected at the afterglow location to a  $2\sigma$  limit of  $f_\nu < 1.5$  mJy (Moin et al. 2009).

<sup>25</sup> <http://www.astronomy.ohio-state.edu/ANDICAM>.

<sup>26</sup> <http://www.astro.yale.edu/smarts>.

### 2.4.3. Spectroscopy

Malesani et al. (2009) obtained a spectrum of the afterglow of GRB 090926A with the X-Shooter instrument mounted on the 8 m Very Large Telescope UT2. Based on the detection of a damped-Ly $\alpha$  system and many strong, narrow absorption features, these authors derive a redshift of 2.1062 for the host galaxy of GRB 090926A.

At this redshift, the prompt isotropic gamma-ray energy release in the  $1\text{--}10^4$  keV bandpass is  $E_{\gamma,\text{iso}} = (1.89 \pm 0.03) \times 10^{54}$  erg. Assuming a nonthermal spectrum up to  $E_{\text{obs}} = 20$  GeV, we infer a lower limit for the initial Lorentz factor of  $\Gamma_0 \gtrsim 700$  (Lithwick & Sari 2001).

## 3. AFTERGLOW MODELING AND RESULTS

In the standard “fireball” model of GRBs (e.g., Piran 2005), a compact central engine (likely either a black hole or a rapidly spinning proto-neutron star) drives a collimated ( $\theta \lesssim 10^\circ$ ), ultrarelativistic ( $\Gamma_0 \gtrsim 100$ ) outflow of matter and/or radiation. Some dissipative process within the outflow (possibly collisionless shocks) gives rise to the prompt gamma-ray emission, converting some fraction ( $\eta_\gamma \equiv E_{\gamma,\text{iso}}/(E_{\gamma,\text{iso}} + E_{\text{KE,iso}})$ ) of the total relativistic energy to high-energy radiation.

The outflow is ultimately slowed as it sweeps up and shock heats the circumburst medium, and synchrotron radiation from electrons accelerated at the shock front results in the broadband afterglow. The resulting spectrum is well described as a series of broken power laws with three characteristic frequencies:  $\nu_a$ , the frequency below which the radiation is self-absorbed;  $\nu_m$ , the characteristic frequency of the emitting electrons; and  $\nu_c$ , the frequency above which electrons are able to cool efficiently through radiation (e.g., Granot & Sari 2002).

The temporal evolution of the afterglow depends on the density profile of the circumburst medium. We consider here two possibilities: a constant-density circumburst medium [ $\rho(r) \propto r^0$ ], as would be expected in an environment similar to the interstellar medium (ISM: Sari et al. 1998), and a wind-like environment [ $\rho(r) \propto r^{-2}$ ], as would be the case for a massive-star progenitor that shed its outer envelope at a constant rate (Chevalier & Li 2000).

At early times, the afterglow emission appears isotropic to distant observers due to the effects of relativistic beaming. However, the outflow slows as it sweeps up more and more circumburst material. When  $\Gamma(t) \approx 1/\theta$ , lateral spreading of the jet becomes important and observers will notice “missing” emission from wide angles (Rhoads 1999; Sari et al. 1999). This hydrodynamic transition manifests itself as an achromatic steepening in the afterglow light curve. Measuring the time of this jet break ( $t_j$ ) allows us to infer the opening angle of the outflow.

In order to ascertain the total relativistic energy output from a GRB, we require three measurements: (1)  $E_{\gamma,\text{iso}}$ , the isotropic energy release in the prompt gamma-ray emission, which is inferred from the high-energy fluence and the associated afterglow or host redshift; (2)  $\theta$ , the half-opening angle of the beamed emission, inferred from the detection of a jet break; and (3)  $E_{\text{KE}}$ , the kinetic energy of the blast wave that is powering the broadband afterglow, which can be inferred either via afterglow modeling or, more accurately, from late-time ra-

dio calorimetry in the nonrelativistic phase (Berger et al. 2004; Frail et al. 2005; van der Horst et al. 2008). We stress here that we are neglecting contributions from non-electromagnetic phenomena (neutrinos, gravity waves, etc.) and slower-moving material (i.e., supernova emission), and so are providing only lower limits on the total energy budget.

Unfortunately, none of the radio afterglows in our sample of LAT events are sufficiently bright to perform calorimetry in the nonrelativistic phase. Instead, we construct afterglow models (including both the standard formulation and corrections for radiative losses and inverse-Compton emission; Sari & Esin 2001) and compare these with our observations using the multi-parameter fitting program of Yost et al. (2003). Our objective is to translate the observed three critical frequencies, together with the peak flux density,  $F_{\nu,\text{max}}$ , and the jet break time,  $t_j$ , into a physical description of the outflow. In particular, we shall attempt to estimate seven parameters:  $E_{\text{KE}}$ , the kinetic energy of the blast wave;  $n$ , the particle density of the circumburst medium (or, alternatively for a wind-like circumburst medium,  $A_*$ , where  $\rho = 5 \times 10^{11} A_* r^{-2} \text{ g cm}^{-3}$ )<sup>27</sup>;  $\epsilon_e$ , the fraction of the total energy apportioned to electrons;  $\epsilon_B$ , the fraction of the total energy apportioned to the magnetic field;  $p$ , the electron power-law index;  $A_V$ , the host-galaxy extinction<sup>28</sup>; and  $\theta$ , the jet half-opening angle.

Optical magnitudes have been converted to flux densities after correcting for Galactic extinction using zero points from Fukugita et al. (1995). To account for differences in instrumental configurations, we have applied a 7% cross-calibration uncertainty to all data points before calculating the models. All reported uncertainties have been determined using a Monte-Carlo bootstrap analysis with 1000 trials and represent only statistical errors associated with the fit. Systematic errors associated with model uncertainties are potentially larger and difficult to estimate.

### 3.1. GRB 090323

#### 3.1.1. Preliminary Considerations

Before proceeding to a detailed model, we derive some initial constraints by looking at the spectral and temporal behavior of the afterglow. Considering first the X-ray afterglow, we fit the flux density to a power-law decay and find a best-fit index of  $\alpha_X = 1.5 \pm 0.1$  ( $\chi^2 = 28.2$  for 22 degrees of freedom, d.o.f.). Performing an analogous power-law fit to the X-ray spectrum, we find  $\beta_X = 1.06^{+0.34}_{-0.13}$  ( $\chi^2 = 8.0$  for 13 d.o.f.; see the on-line compilation of N.R.B. for details).

We perform a similar fit in the optical ( $r'$ ,  $R$ , and  $i'$  filters), forcing the decay index to be identical in all filters and ignoring the points at  $t > 10$  days due to host galaxy contamination (see below). Though the fit quality

<sup>27</sup> For a wind-like medium, the circumburst density is normalized for a progenitor mass-loss rate of  $\dot{M} = 10^{-5} \text{ M}_\odot \text{ yr}^{-1}$  and a wind speed of  $v_w = 1000 \text{ km s}^{-1}$  (Chevalier & Li 1999). The units of  $A_*$  are thus  $\text{g cm}^{-1}$ , and the conversion to a particle density (as a function of radius) is given by  $n = 30 A_* r_{17}^{-2} \text{ cm}^{-3}$ .

<sup>28</sup> We have assumed an SMC-like extinction curve for all events here (Pei 1992; Kann et al. 2006). Given the relatively modest amounts of dust inferred for these host galaxies, this choice does not affect any of our primary conclusions.

TABLE 4  
GRB 090323 AFTERGLOW BEST-FIT PARAMETERS

$E_{\text{KE,iso}}$ ( $10^{52}$ erg)	$A_*$ ( $\text{g cm}^{-1}$ )	$\epsilon_e$ (%)	$\epsilon_B$ (%)	$\theta$ ( $^\circ$ )	$p$	$A_V(\text{host})$ (mag)	$\chi^2_\nu$ (d.o.f.)
$200^{+90}_{-30}$	$0.12^{+0.02}_{-0.01}$	$6.8 \pm 0.5$	$0.31 \pm 0.12$	$2.6^{+0.6}_{-0.1}$	$2.81 \pm 0.06$	$0.12 \pm 0.03$	1.39 (70)

is much worse ( $\chi^2 = 178.4$  for 34 d.o.f.), we find a similar decay index as seen in the X-rays:  $\alpha_O = 1.8 \pm 0.1$ . Likewise, a spectral fit of the multi-color GROND data obtained at  $t = 1.12$  days results in a spectral index of  $\beta_O = 0.95 \pm 0.01$ .

The comparable spectral and temporal indices in the optical and X-rays suggests that both bandpasses fall in the same synchrotron spectral regime. In other words, the cooling frequency should fall either below the optical or above the X-ray bandpass for the duration of these observations ( $t \approx 1\text{--}10$  days). Examining the “closure” relations between  $\alpha$  and  $\beta$  in different circumburst media and spectral regimes (e.g., Price et al. 2002), we find that we can rule out a low cooling frequency at large significance, as this would require  $\alpha = (3\beta - 1)/2 \approx 1.0$  for both a wind-like and ISM-like circumburst medium.

The relatively flat radio light curve (modulo effects from interstellar scintillation) until  $t \approx 20$  days has two important implications. First, it suggests a wind-like circumburst medium, as the flux would be expected to rise in proportion to  $t^{1/2}$  for  $\nu_a < \nu < \nu_m$  in a constant-density environment. From this and our previous X-ray and optical spectral and temporal decay indices, we can infer that the electron spectral index is relatively steep,  $p \approx 2.7$  (e.g., Shen et al. 2006; Starling et al. 2008; Curran et al. 2009). Second, together with the lack of evidence for late-time steepening in the X-ray and optical light curves, this suggests that if a jet break is required, it should occur at  $t \gtrsim 20$  days. (Another possibility for the decay could be the peak frequency  $\nu_m$  passing through the radio bands.)

Finally, the relatively bright  $r'$ -band detection at  $t = 130$  days is almost certainly dominated by flux from the host galaxy and not the afterglow of GRB 090323. If we assume this flux is due entirely to host light, the host will contribute a significant fraction of the flux in some of the late-time points of our optical light curve ( $\sim 40\%$  at  $t = 14$  days in  $r'$ ). We assume a flat spectrum and subtract a host contribution of  $f_\nu = 0.40 \mu\text{Jy}$  from all optical data points for our afterglow modeling.

### 3.1.2. Modeling Results

With the above constraints in hand, we have modeled the afterglow of GRB 090323 with the software described above. The resulting best-fit wind model is plotted in Figure 2 and the derived parameters are provided in Table 4. The overall fit quality is reasonable ( $\chi^2 = 97.4$  for 70 d.o.f.), with comparable residuals in the radio, optical, and X-ray bandpasses. We were unable to find any physically plausible models for a constant-density medium with reasonable fit quality.

As expected, the cooling frequency  $\nu_c$  lies at or above the X-ray bandpass for the duration of our observations ( $t \gtrsim 1$  day). As a result, the ratio of  $E_{\text{KE,iso}}$  to  $A_*$  is required to be larger than inferred for previ-

ous GRBs (e.g., Panaitescu & Kumar 2001b; Yost et al. 2003). While the isotropic blast-wave kinetic energy is relatively large compared to that of previously modeled GRBs ( $E_{\text{KE,iso}} = 2.0 \times 10^{54}$  erg), it is in fact comparable to the prompt gamma-ray energy release, as would be expected for reasonable values of the gamma-ray efficiency  $\eta_\gamma$ . The inferred density is slightly lower than usual ( $A_* = 0.12 \text{ g cm}^{-1}$ ), though smaller values have been reported in the literature (e.g., GRB 020405; Chevalier et al. 2004).

The models result in a jet break time of  $t_j = 17.8^{+19.6}_{-1.7}$  days. This occurs after the X-ray and optical observations have stopped (except for the host detection in the optical), and is therefore most directly constrained by the radio light curve. In Figure 2 we also plot the same model parameters but for an isotropic explosion, where the turnover in the radio at late times is due to the peak frequency  $\nu_m$  passing through the radio bandpass. The current radio data provide only weak constraints on the jet break time upper bound, although the models suggest a much smaller uncertainty in the opening angle. We return to the robustness of our determination of  $\theta$  in § 4.4.

While the jet break occurs rather late in the observer frame (e.g., Zeh et al. 2006), the inferred opening angle ( $\theta = 2.6^\circ$ ) is still relatively small compared with previous samples. Though the dependence is not strong ( $\theta \propto (E_{\text{KE,iso}}/A_*)^{-1/4}$ ), the large ratio of  $E_{\text{KE,iso}}$  to  $A_*$  effectively lowers the opening angle for a given jet break time.

After applying the collimation correction, we find that the true energy release of GRB 090323 is  $E_\gamma = 4.1^{+2.9}_{-0.8} \times 10^{51}$  erg,  $E_{\text{KE}} = 2.1^{+2.4}_{-0.5} \times 10^{51}$  erg. We compare these results with a larger sample of events in § 4.1.

## 3.2. GRB 090328

### 3.2.1. Preliminary Considerations

Following the results of § 3.1.1, we first consider the spectral and temporal indices of GRB 090328 in the observed bandpasses independently. The X-ray afterglow light curve is well fit by a single power-law decay with index  $\alpha_X = 1.65 \pm 0.09$  ( $\chi^2 = 24.1$  for 26 d.o.f.). The spectral index, on the other hand, is less well constrained:  $\beta_X = 1.1^{+0.4}_{-0.3}$  ( $\chi^2 = 6.7$  for 14 d.o.f.). We note that the X-ray spectral fit indicates the presence of an additional absorption component in the host galaxy of GRB 090328 ( $N_H = 4.9^{+3.5}_{-2.8} \times 10^{21} \text{ cm}^{-2}$  at  $z = 0.736$ ).

The relative paucity of optical data, particularly at late times, makes a similar analysis challenging. Fitting the UVOT  $U$ -band light curve to a single power-law decay results in an index of  $\alpha_O = 1.2 \pm 0.1$ ; however, the quality of the fit is extremely poor ( $\chi^2 = 24.5$  for 5 d.o.f.). An initially steep decay in the light curve becomes more shallow at later times, much like is seen in the UVOT

TABLE 5  
GRB 090328 AFTERGLOW BEST-FIT PARAMETERS

$E_{\text{KE,iso}}$ ( $10^{52}$ erg)	$A_*$ (g cm $^{-1}$ )	$\epsilon_e$ (%)	$\epsilon_B$ (%)	$\theta$ ( $^\circ$ )	$p$	$A_V(\text{host})$ (mag)	$\chi^2_\nu$ (d.o.f.)
$11^{+5}_{-3}$	$0.11^{+0.03}_{-0.05}$	$9 \pm 2$	$0.9^{+0.7}_{-0.2}$	$5.2^{+1.4}_{-0.7}$	$2.58^{+0.12}_{-0.07}$	$0.41 \pm 0.04$	1.63 (43)

white filter (Marshall et al. 2009). This result suggests the presence of host-galaxy contamination at late times.

The best-fit spectral index assuming a power-law model for the simultaneous multi-color GROND data is  $\beta_O = 1.24 \pm 0.04$ , although the fit quality is extremely poor ( $\chi^2 = 85.5$  for 5 d.o.f.). The poor fit statistic is dominated almost entirely by a single filter (*J* band), however, and the relatively steep spectral index (compared to  $\beta_{OX} \approx 0.8$ ) appears to be robust. This is consistent with the presence of dust indicated by the X-ray spectral fits.

In the case of GRB 090328, examining the  $\alpha$ - $\beta$  closure relations yields little insight. Because of the poor fit quality in the optical and the large uncertainty in the X-ray spectral index, we are unable to rule out much of the available parameter space. We note, however, that with the exception of the excess absorption, the X-ray afterglow light curve and spectrum of GRB 090328 closely resemble those of GRB 090323. Similarly, if we assume that the late-time optical data are dominated by host-galaxy light, the optical light curve is also compatible with that seen from GRB 090323.

### 3.2.2. Modeling Results

The results of our best-fit model for the afterglow of GRB 090328 are shown in Figure 4 and the parameters are provided in Table 5. The overall fit quality is reasonable ( $\chi^2 = 70.1$  for 43 d.o.f.). Only the early radio light curve, which suffers from strong scintillation, differs systematically from the model predictions.

Much like in the case of GRB 090323, we find that only a wind-like circumburst medium can reproduce the observed light curves with plausible physical parameters. While the slow rise in the early radio afterglow is better explained by a constant-density environment, our modeling indicates ISM-like solutions require  $\epsilon_e \rightarrow 1$  and  $\epsilon_B \rightarrow 0$  (i.e., a fully radiative solution).

We note that our solutions require the presence of a relatively bright host galaxy — if the late-time *U*-band points are not predominantly due to host light, the fit quality of our models decreases somewhat ( $\chi^2_\nu \gtrsim 2.0$ ).

The steep decline in the radio light curve at late times offers relatively tight constraints on the presence of a jet break. In fact, the break is consistent with (though not formally required by) the final few points in the X-ray light curve as well. For our best-fit model, we infer  $t_j = 6.4^{+12.0}_{-1.5}$  days and  $\theta = (5.2^{+1.4}_{-0.7})^\circ$ . Primarily because of the relatively sparse optical data, we find several unique models with similar though slightly worse overall fit statistics. We note, however, that all these solutions result in comparable opening angles, the most important parameter for our energetics calculations.

Correcting for the effects of beaming, we find an energy release of  $E_\gamma = 5.5^{+4.3}_{-1.8} \times 10^{50}$  erg and  $E_{\text{KE}} = 4.5^{+6.1}_{-2.0} \times 10^{50}$  erg. The total relativistic energy release,  $E_{\text{tot}} \approx$

$10^{51}$  erg, is somewhat less than the value we derive for GRB 090323.

### 3.3. GRB 090902B

#### 3.3.1. Preliminary Considerations

A single power-law model with  $\alpha_X = 1.36 \pm 0.03$  provides a good fit to the X-ray light curve of GRB 090902B over the entire span of the XRT observations ( $\chi^2 = 94.8$  for 107 d.o.f.). Likewise, the X-ray spectrum is well fit by a power law with  $\beta_X = 0.90 \pm 0.13$  ( $\chi^2 = 42.8$  for 56 d.o.f.). The X-ray spectrum strongly favors the presence of absorption in excess of the Galactic value along the GRB line of sight. Assuming the dust arises in the GRB host galaxy at  $z = 1.82$ , the best-fit host column density is  $N_H = (6.3^{+2.1}_{-1.7}) \times 10^{21}$  cm $^{-2}$ .

The decline inferred from the early-time ROTSE-III *R*-band observations is significantly steeper than the late-time ( $t \gtrsim 1$  day) optical decay. Following Pandey et al. (2010), we attribute this emission largely to the presence of a reverse shock and do not include these data points in any subsequent analysis. Simultaneously fitting the remaining optical data to a power law of the form  $f_\nu \propto t^{-\alpha} \nu^{-\beta}$ , we find  $\alpha_O = 0.89 \pm 0.05$  and  $\beta_O = 0.76 \pm 0.07$  ( $\chi^2 = 36.4$  for 18 d.o.f.).

The large discrepancy between the X-ray and optical temporal decay slopes suggests that the two bandpasses fall in different synchrotron spectral regimes (i.e.,  $\nu_O < \nu_c < \nu_X$ ). Furthermore, the relatively shallow optical decline strongly rules out a wind-like circumburst medium. We find that both the X-ray and optical bandpasses are broadly consistent with expansion into a constant-density medium and an electron index  $p \approx 2.2$ . The observed X-ray temporal decline is somewhat steeper than what is predicted, but this could be accounted for by synchrotron radiative losses at later times.

The radio light curve appears to decay from the earliest observations at  $t_0 + 1.5$  days onward. This behavior is not expected in the simple forward-shock model where radio afterglows are typically observed to rise (for a constant-density medium) or remain flat (for a stratified environment) until either a jet break occurs or the peak frequency reaches the radio bandpass (typically weeks after the burst).

However, bright, early-time radio emission is actually rather common, having been detected in a quarter of all radio afterglows to date (Soderberg & Ramirez-Ruiz 2003; Chandra et al. 2010). This prompt component is most commonly attributed to the reverse shock propagating in the shocked ejecta. The reverse-shock emission fades away quite quickly, and at late times ( $t \gtrsim 5$  days) the afterglow will be dominated by the forward-shock emission. GRB 990123 was the first known example (Kulkarni et al. 1999), but there have been many more claimed detections since (e.g., Berger et al. 2003b; Panaitescu 2005; Chandra et al. 2010).



TABLE 6  
GRB 090902B AFTERGLOW BEST-FIT PARAMETERS

$E_{\text{KE,iso}}$ ( $10^{52}$ erg)	$n$ ( $\text{cm}^{-3}$ )	$\epsilon_e$ (%)	$\epsilon_B$ (%)	$\theta$ ( $^\circ$ )	$p$	$A_V(\text{host})$ (mag)	$\chi^2_{\nu}$ (d.o.f.)
$68^{+14}_{-6}$	$(5.8^{+1.2}_{-1.8}) \times 10^{-4}$	$15^{+4}_{-1}$	$5.8^{+2.5}_{-1.0}$	$3.4^{+0.4}_{-0.3}$	$2.22^{+0.08}_{-0.04}$	$0.18^{+0.08}_{-0.05}$	1.06 (137)

The radio light curve of GRB 090902B most closely resembles that of GRB 991216 in which the early power-law decline is due to the overlap of emission from the reverse and forward shocks (Frail et al. 2000b). Since we are concerned here with the modeling of the behavior of the forward shock, we will exclude the early radio point and postpone a full discussion to a subsequent paper dealing with a compilation all known GRBs with prompt radio emission (Chandra et al. 2010).

### 3.3.2. Modeling Results

Using the constraints derived above, our best-fit model for GRB 090902B assuming a constant-density circumburst medium is shown in Figure 6, and the derived model parameters are presented in Table 6. The overall fit quality is quite reasonable ( $\chi^2 = 145.8$  for 137 d.o.f.), although the fit quality is somewhat better in the X-ray and radio than in the optical/UV. We stress, however, that we have not included the first two *R*-band data points and the first VLA detection in our fitting, as we believe the flux at these points is dominated by reverse-shock emission.

Much like the other events considered here, our inferred parameters for GRB 090902B suggest a large afterglow kinetic energy ( $E_{\text{KE,iso}} = 6.8^{+1.4}_{-0.6} \times 10^{53}$  erg; a factor of five less than the prompt gamma-ray energy release) and a low circumburst density ( $n = 5.8^{+1.2}_{-1.8} \times 10^{-4} \text{ cm}^{-3}$ ). In fact, we are unable to find any acceptable solutions ( $\chi^2_{\nu} < 2$ ) with  $n > 10^{-2} \text{ cm}^{-3}$ . The inferred circumburst density for many long-duration GRBs is lower than typical values observed in dense molecular clouds (where presumably their massive-star progenitors have formed). Yet the value we have derived for GRB 090902B is closer in fact to what one might find in the ambient ISM or even the intergalactic medium (IGM), as has been found for the few short-hard GRBs with sufficient afterglow data (e.g., Panaitescu 2006; Perley et al. 2009b). We return to this issue in more detail in § 4.3.

Neither the X-ray nor the (less constraining) optical bandpass formally require a jet break over the duration of our observations. In this case, like GRB 090323, we must rely largely on the radio. Our models suggest that the peak synchrotron frequency,  $\nu_m$ , will not fall to the radio bandpass until  $t \gtrsim 40$  days. The declining radio light curve at late times therefore requires a jet break at  $t_j = 6.2^{+2.4}_{-0.8}$  days. Again, because of the large ratio of  $E_{\text{KE,iso}}$  to  $n$ , this corresponds to a relatively narrow opening angle:  $\theta = (3.4^{+0.4}_{-0.3})^\circ$ .

Correcting for collimation, we find the true energy release from GRB 090902B to be  $E_{\gamma} = (5.6 \pm 1.5) \times 10^{51}$  erg,  $E_{\text{KE}} = 1.2^{+0.7}_{-0.4} \times 10^{51}$  erg.

### 3.4. GRB 090926A

#### 3.4.1. Preliminary Considerations

Fitting a single power-law decay to the X-ray light curve, we find a reasonable quality fit ( $\chi^2 = 120.8$  for 97 d.o.f.) with  $\alpha_X = 1.43 \pm 0.03$ . The fit quality improves somewhat if we allow for a steeper decay at late times, although the post-break decay index is not particularly well constrained:  $\alpha_{X,1} = 1.35 \pm 0.04$ ,  $\alpha_{X,2} = 2.6^{+0.9}_{-0.5}$ ,  $t_b = 9.1^{+2.1}_{-1.4}$  days,  $\chi^2 = 111.9$  (95 d.o.f.). The X-ray spectrum is well fit ( $\chi^2 = 41.7$  for 51 d.o.f.) by a single power law with index  $\beta_X = 1.12 \pm 0.13$  and does not require any absorption in addition to the Galactic component.

The behavior in the optical bandpass, however, is significantly more complex (Fig. 8). The flux in the initial UVOT *U*- and *V*-band observations declines until  $t \approx 0.8$  day, at which point a rebrightening or plateau phase is evident in all five observed filters (*UBVRI*). The optical light curve of GRB 090926A is reminiscent of the rebrightening seen at  $t \approx 1$  day from GRB 970508 (Djorgovski et al. 1997) and GRB 071003 (Perley et al. 2008 and references therein). Such a phenomenon is difficult to reconcile with the standard afterglow paradigm. Possible explanations include a sharp change in the circumburst density (e.g., Lazzati et al. 2002; Tam et al. 2005 but, c.f. Nakar & Granot 2007) or a smooth injection of energy into the forward shock from the central engine (e.g., Rees & Meszaros 1998). A more detailed analysis of the early-time behavior of GRB 090926A is beyond the scope of this work.

The optical light curve appears to peak at  $t \approx 1$  day in all bands. Fitting the PROMPT *BVRI* data at  $t > 1$  day to a single power law results in a best-fit index of  $\alpha_O = 1.38 \pm 0.02$  ( $\chi^2 = 181.6$  for 119 d.o.f.). However, extrapolating these results to the late-time Gemini imaging at  $t \approx 23$  days greatly overestimates the observed flux (Fig. 9). Much like the X-ray data, this strongly suggests a steepening of the optical light curve at  $t \gtrsim 8$  d.

We have also performed a joint spectral and temporal fit to the PROMPT data to estimate the optical spectral index,  $\beta_O$ . We include only the *V*-, *R*-, and *I*-band data, as both the *U* and *B* bands are likely affected by Ly $\alpha$  absorption at  $z = 2.11$ . We then find a best-fit optical spectral index of  $\beta_O = 1.03 \pm 0.05$ . The formal fit quality is relatively poor, however ( $\chi^2 = 183.2$  for 68 d.o.f.). Finally, the optical to X-ray spectral index at  $t \gtrsim 1$  day is  $\beta_{OX} \approx 1.1$ .

Taken together, we find similar temporal ( $\alpha_{X,1} = 1.35$ ,  $\alpha_{O,1} = 1.38$ ) and spectral ( $\beta_X = 1.12$ ,  $\beta_O = 1.03$ ) indices in the X-ray and optical for  $t \approx 1$ –9 days. Furthermore, the optical to X-ray spectral index,  $\beta_{OX}$ , is comparable to both the optical and X-ray spectral indices. These facts strongly suggest that both the X-ray and optical bandpasses fall in the same synchrotron spectral regime.

Considering the various synchrotron closure relations, the best fit appears to occur when both bandpasses fall above the cooling frequency,  $\nu_c$ . The afterglow decay in

TABLE 7  
GRB 090926A AFTERGLOW BEST-FIT PARAMETERS

$E_{\text{KE,iso}}$ ( $10^{52}$ erg)	$A_*$ ( $\text{g cm}^{-1}$ )	$\epsilon_e$ (%)	$\epsilon_B$ (%)	$\theta$ ( $^\circ$ )	$p$	$A_V(\text{host})$ (mag)	$\chi^2_\nu$ (d.o.f.)
$24.2^{+1.9}_{-1.3}$	$0.67^{+0.14}_{-0.02}$	$30^{+2}_{-3}$	$0.87^{+0.07}_{-0.03}$	$7.1 \pm 0.2$	$2.19 \pm 0.02$	$0.06 \pm 0.01$	1.18 (149)

this regime is independent of circumburst medium. Such a low cooling frequency is somewhat unusual, though not unprecedented (e.g., GRB 050904; Frail et al. 2006) in GRB afterglows. The implied electron index in this case would be  $p \approx 2.3$ .

Finally, we note that the steepening in both the X-ray and optical bands at  $t \approx 9$  days is strongly suggestive of a jet break. We examine this possibility in greater detail in the following section.

### 3.4.2. Modeling Results

Unfortunately, without a radio light curve we cannot uniquely solve for the physical parameters of the afterglow of GRB 090926A. We can, however, at the very least more robustly constrain the jet break time and opening angle of this event, as well as identify broad trends in the parameters associated with acceptable solutions. As discussed previously, we consider only data at  $t \geq 1$  day due to the rebrightening observed in the optical before then.

As expected, we were able to find a number of acceptable solutions, both for constant-density and wind-like environments. One example for a wind-like circumburst environment is shown in Table 7 and Figure 9. The model provides a reasonable description of the data in all bandpasses, with  $\chi^2 = 175.1$  (149 d.o.f.). Without radio data, however, the solution presented in Table 7 is not unique in a global sense, and therefore should only be taken as representative of a larger family of solutions.

Most importantly for our purposes, a jet break at  $t \approx 9 \pm 2$  days is required to provide a reasonable fit to the late-time X-ray and (particularly) optical data (Fig. 9). The precise translation into an opening angle is dependent on the circumburst density (both profile and absolute value) and the afterglow energy. Taking the values derived above into account, we find a typical value for a wind-like circumburst medium of  $\theta \approx 7^\circ$ . Based on the observed spread of  $E_{\text{KE}}$  and  $A_*$  in our wind models, we adopt an approximate uncertainty in this value of  $\theta \approx (7^{+3}_{-1})^\circ$ .

Correcting for collimation, we find a prompt energy release of  $E_\gamma = 1.4^{+1.5}_{-0.4} \times 10^{52}$  erg. Assuming a reasonable value for the gamma-ray efficiency ( $\eta_\gamma \gtrsim 10\%$ ), we adopt a lower limit on the isotropic blast-wave energy of  $E_{\text{KE,iso}} \gtrsim 10^{53}$  erg. We then find a collimation-corrected afterglow energy of  $E_{\text{KE}} \gtrsim 5 \times 10^{50}$  erg.

## 4. DISCUSSION

### 4.1. Central-Engine Constraints I: Energetics and Remnants

In Table 8, we summarize the primary results from this work, including the redshift, initial Lorentz factor, beaming angle, density, and collimation-corrected energy release for each of the four LAT GRBs considered here.

Before the launch of the *Swift* satellite in 2004, the majority of all well-observed afterglows were inferred to be highly collimated, with opening angles  $\theta \lesssim 10^\circ$  (Zeh et al. 2006). The most notable exceptions were the handful of the most nearby events, including GRB 980425 (Galama et al. 1998; Kulkarni et al. 1998), GRB 031203 (Soderberg et al. 2004; Sazonov et al. 2004; Thomsen et al. 2004; Cobb et al. 2004; Malesani et al. 2004; Gal-Yam et al. 2004), and (post-*Swift*) GRB 060218 (Mirabal et al. 2006; Campana et al. 2006; Soderberg et al. 2006; Modjaz et al. 2006; Sollerman et al. 2006; Pian et al. 2006; Ferrero et al. 2006; Maeda et al. 2007), all of which appear to be isotropic explosions that were energetically dominated by their nonrelativistic ejecta (i.e., their associated supernovae).

The typical GRB afterglows discovered by *Swift*, however, did not fit neatly into this simple bimodal picture. The afterglows of most *Swift* GRBs, including the large fraction at cosmological distances, exhibit a much broader range of opening angles (Kocevski & Butler 2008; Racusin et al. 2009), with some extreme events lacking a detectable jet break signature in the X-rays out to hundreds of days after the burst (e.g., GRB 060729; Grupe et al. 2007, 2010).

All four of the LAT-detected events we have studied here are consistent with a relatively high degree of collimation ( $\theta \lesssim 10^\circ$ ). In this respect, then, the afterglows of LAT-detected GRBs more closely resemble the pre-*Swift* sample. This in and of itself is not entirely surprising, given the tremendous high- $E_{\gamma,\text{iso}}$  bias for events detected by the LAT (Figure 1).

In Figure 10, we plot the two-dimensional (prompt + afterglow) collimation-corrected relativistic energy release for the four LAT events in this work, compared with several additional samples. Shown in red are 11 pre-*Swift* GRBs at cosmological distances ( $z > 0.5$ ) for which sufficient broadband (X-ray, optical, and radio) data exist to constrain both  $E_\gamma$  and  $E_{\text{KE}}$ . We find the logarithmic mean for the sum of these two values,  $E_{\text{rel}} \approx E_\gamma + E_{\text{KE}}$ ,<sup>29</sup> to be  $\langle E_{\text{rel}} \rangle = 2.0 \times 10^{51}$  erg, with an error in the mean of 0.17 dex. The solid black line indicates a constant value of  $E_{\text{rel}}$  corresponding to this mean, while the gray shaded regions indicate  $1\sigma$ ,  $2\sigma$ , and  $3\sigma$  confidence intervals.

The gray dotted line in Figure 10 represents a constant gamma-ray efficiency of  $\eta_\gamma = 50\%$ . Most events are roughly consistent (or even exceed) this value. This result presents a problem for most internal shock models of the prompt emission, which predict a maximal gamma-ray efficiency of  $\eta_\gamma \lesssim 10\%$  (Kobayashi et al. 1997; Daigne & Mochkovitch 1998). Alternatively, if the

<sup>29</sup> Assuming an approximately log-normal distribution, we calculate the weighted mean of  $\langle \log_{10}(E_{\text{tot}}) \rangle$ , where  $\sigma = \delta E_{\text{tot}} / (\ln(10) \times E_{\text{tot}})$ .

TABLE 8  
SUMMARY OF GRB PARAMETERS

GRB	$z$	$f_{\gamma}^a$ ( $10^{-4}$ erg cm $^{-2}$ )	$E_{\gamma, \text{iso}}^a$ ( $10^{51}$ erg)	$\Gamma_0$	$\theta$ ( $^{\circ}$ )	$E_{\gamma}$ ( $10^{51}$ erg)	$E_{\text{KE}}$ ( $10^{51}$ erg)	$n/A_*$ (cm $^{-3}$ / g cm $^{-1}$ )
090323	3.568	$1.50 \pm 0.20$	$3.99 \pm 0.53$	$\gtrsim 600$	$2.6^{+0.6}_{-0.1}$	$4.1^{+2.9}_{-0.8}$	$2.1^{+2.4}_{-0.5}$	$0.12^{+0.02}_{-0.01}$
090328	0.7357	$0.73 \pm 0.08$	$0.134 \pm 0.014$	$\gtrsim 200$	$5.2^{+1.4}_{-0.7}$	$0.55^{+0.43}_{-0.18}$	$0.45^{+0.61}_{-0.20}$	$0.11^{+0.03}_{-0.05}$
090902B	1.8229	$3.83 \pm 0.05$	$3.20 \pm 0.04$	$\gtrsim 1000$	$3.4^{+0.4}_{-0.3}$	$5.6 \pm 1.5$	$1.2^{+0.7}_{-0.4}$	$(5.8^{+1.2}_{-1.8}) \times 10^{-4}$
090926A	2.1062	$1.73 \pm 0.03$	$1.89 \pm 0.03$	$\gtrsim 700$	$7^{+3}_{-1}$	$14^{+15}_{-4}$	$\gtrsim 0.5$	$\dots^b$

<sup>a</sup> 1–10<sup>4</sup> keV observer frame bandpass.

<sup>b</sup> We have not included an estimate for the density of GRB 090926A, as this parameter was not well constrained due to the lack of radio data.

early afterglow undergoes a relatively long-lived radiative phase, we may be significantly underestimating the blastwave kinetic energy (Ghisellini et al. 2010).

As discussed previously, the three most nearby GRBs are clearly subenergetic outliers. The energy release from the supernovae associated with these events,  $E_{\text{SN}} \sim 10^{51}$ – $10^{52}$  erg (Iwamoto et al. 1998; Mazzali et al. 2006), is several orders of magnitude larger than the energy apportioned to the relativistic ejecta. To further distinguish these events, the rate of such subluminal GRBs suggests they are many times more common (per unit volume) than the typical cosmological GRBs (Soderberg et al. 2006; Cobb et al. 2006; Guetta & Valle 2007), although Butler et al. (2010) have suggested that both samples may be described by a single luminosity function — that is, subluminal GRBs may simply extend continuously from the higher-energy population.

Of the LAT events studied here, the total relativistic energy output from only a single one (GRB 090328) is consistent with the pre-*Swift* mean at the  $3\sigma$  level. This is not entirely surprising, however, as the  $E_{\text{rel}}$  distribution of pre-*Swift* GRBs exhibits a reasonable dispersion ( $\sim 0.55$  dex, or a factor of 3.5). It is clear, however, that GRB 090323, GRB 090902B, and GRB 090926A all fall at the very high end of the pre-*Swift* distribution. In particular, the only event comparable to GRB 090926A, with  $E_{\gamma} \approx 10^{52}$  erg and  $E_{\text{KE}}$  relatively unconstrained, is GRB 970508 ( $E_{\text{rel}} \approx 1.5 \times 10^{52}$  erg, dominated by the large afterglow kinetic energy; Yost et al. 2003; Berger et al. 2004).

GRBs 090323, 090902B, and 090926A appear consistent instead with a subsample of the brightest *Swift* events from Cenko et al. (2010) (see also Frail et al. 2006; Chandra et al. 2008). This, too, is not unexpected, as these *Swift* events were chosen on the basis of large  $E_{\gamma, \text{iso}}$  values, and therefore should in large part mimic at least this component of the LAT selection effects.

Much like several events in the bright *Swift* sample, GRB 090926A appears to exceed the canonical GRB relativistic energy release of  $10^{51}$  erg by roughly an order of magnitude. We stress that our methodology provides a relatively conservative estimate for the prompt energy release of GRB 090926A, for a number of reasons. First, we consider only the rest-frame 1–10<sup>4</sup> keV bandpass for *all* events considered in Figure 10. Extrapolating measurements from previous instruments, with bandpasses typically extending only to  $\sim 1$  MeV (only a few hundred keV for *Swift*), up to the GeV range would introduce significant uncertainties in  $E_{\gamma, \text{iso}}$ . While allowing for a more robust burst-to-burst comparison, we have not included

a significant fraction of the detected high-energy fluence in our energy calculations for these LAT events ( $\sim 30\%$  for GRB 090926A).

Second, we have derived the opening angles of all our events using parameters inferred from our broadband modeling. Consequently, the  $E_{\text{rel}}$  we derive for GRB 090902B is a factor of several lower than that reported by other authors (§ 4.3 and 4.4). Most importantly, for GRB 090926A we have deliberately used only wind-like models to derive the beaming angle, even though our data are insufficient to rule out a constant-density medium. For ISM-like models, the opening angles we found were on average  $\sim 75\%$  larger, corresponding to a beaming-corrected prompt energy release of  $E_{\gamma} \approx 4 \times 10^{52}$  erg.

Together with the discovery of several comparable events to GRB 090926A in the past few years, including GRB 050904 ( $E_{\text{rel}} \approx 2 \times 10^{52}$  erg; Frail et al. 2006), GRB 070125 ( $E_{\text{rel}} \approx 3 \times 10^{52}$  erg; Chandra et al. 2008), GRB 050820A ( $E_{\text{rel}} \approx 4 \times 10^{52}$  erg; Cenko et al. 2010), and GRB 090423 ( $E_{\text{rel}} \gtrsim 1 \times 10^{52}$  erg; Chandra et al. 2010), we now believe there is substantial evidence in favor of a subpopulation of GRBs with relativistic energy outputs either very near or above  $10^{52}$  erg. We refer to such events as hyper-energetic GRBs in what follows, and outline some of the implications of this particular energy threshold.

Much as was argued by Starling et al. (2009) for the case of GRB 080721, the total energy budget is an important diagnostic for any central-engine model. In particular, models for which the outflow is powered by the spin-down of a highly magnetized ( $B \gtrsim 10^{15}$  G) proto-neutron star are subject to strict constraints on the total energy budget of  $E_{\text{total}} < 3 \times 10^{52}$  erg (the rotational energy of a maximally spinning  $1.4 M_{\odot}$  neutron star; e.g., Thompson et al. 2004; Metzger et al. 2007).

Only accounting for the relativistic energy output, GRB 090926A seems to approach within at least a factor of a few of this limit. While we caution that there are still significant uncertainties associated with the models used to infer the afterglow parameters (and hence  $E_{\text{rel}}$ ), we have not yet accounted for additional sources of energy, including (nonrelativistic) SN emission, radiative losses at early times due to bright X-ray flares (Burrows et al. 2005a; Falcone et al. 2007), and synchrotron losses during the later phases of afterglow evolution (Yost et al. 2003). Even if we have overestimated the relativistic energy output for these events by a factor of several and the magnetar energy limit is not strictly violated, the

tremendous efficiency required by this process strains credulity.

#### 4.2. Central-Engine Constraints II: Lorentz Factor, Opening Angle, and Acceleration Mechanisms

Unlike magnetar models that are powered by the spin-down of a highly magnetized proto-neutron star, GRB models in which the core of an evolved massive star collapses promptly to form a black hole and accretion disk system (“Type I collapsars” following the nomenclature of MacFadyen et al. 2001) have significantly relaxed constraints on the total energy budget that are in at least some cases capable of accommodating hyper-energetic events. For instance, the initial models of MacFadyen & Woosley (1999) that begin with a  $35 M_{\odot}$  He star with a  $10 M_{\odot}$  evolved core lead to accretion rates of  $\sim 0.1 M_{\odot} \text{ s}^{-1}$  onto a  $3 M_{\odot}$  rotating black hole. Assuming approximately continuous feeding over the fallback time of the stellar envelope ( $\sim 10$  s), this corresponds to a total accreted mass  $\sim 1 M_{\odot}$ . In addition to the accretion process, an even larger reserve lies in the rotational energy of the black hole:

$$E_{\text{rot}} = \frac{M_{\text{BH}} c^2}{2} \left\{ 2 - \left[ \left( 1 + \sqrt{1 - a^2} \right)^2 + a^2 \right]^{1/2} \right\}, \quad (1)$$

where  $M_{\text{BH}}$  is the black hole mass and  $a$  is the dimensionless rotation parameter ( $a \equiv Jc/M_{\text{BH}}^2 G$ ).

At least two competing theories exist, however, to explain how the system can channel this energy to produce the collimated, relativistic jets we observe from GRBs. First, the energy may be extracted from the accretion process via  $\nu\bar{\nu}$  annihilation (e.g., Woosley 1993; MacFadyen & Woosley 1999; Popham et al. 1999; Ruffert & Janka 1999; Narayan et al. 2001). At the hyper-Eddington rates expected for collapsars, the accretion disk formed around the rotating black hole will be optically thick and photons are unable to escape. The viscous heat is instead balanced by cooling via neutrino emission. Annihilation of  $\nu\bar{\nu}$  pairs will produce a gas of hot electron-positron pairs, which, assuming sufficiently low baryon loading, can then rapidly expand in the low-density regions along the axis of rotation (a “fireball”, e.g., Zhang et al. 2003).

Alternatively, if the black hole is rotating and the accretion disk is threaded by sufficiently large magnetic fields ( $B \approx 10^{15}$  G for a  $3 M_{\odot}$  black hole), energy can be extracted directly from the rotating black hole via the Blandford-Znajek mechanism (e.g., Blandford & Znajek 1977; Lee et al. 2000). The accretion disk serves primarily to anchor the large magnetic field (which would otherwise disperse), and the energy is emitted as a largely Poynting flux-dominated outflow.

The efficiency of converting the potential energy of the rotating black hole plus accretion disk system into a form suitable to launch a collimated, relativistic outflow has been intensively studied in the last decade. In the case of  $\nu\bar{\nu}$  annihilation, the overall efficiency of converting the neutrino luminosity of the cooling disk ( $L_{\nu}$ ) into  $e^+e^-$  pairs ( $L_{\nu\bar{\nu}}$ ) depends sensitively on the mass accretion rate, the disk geometry, and (perhaps) the effects of general relativity on neutrino physics (e.g., Popham et al. 1999; MacFadyen & Woosley 1999;

Ruffert & Janka 1999). But the majority of recent simulations suggest at most a modest efficiency ( $\eta_{\nu\bar{\nu}} \equiv L_{\nu\bar{\nu}}/L_{\nu} \lesssim 0.5\%$ ) for the energy available to launch a relativistic jet (e.g., Rosswog et al. 2003; Birkel et al. 2007; Harikae et al. 2010). For typical disk luminosities of  $L_{\nu} \approx 10^{51}\text{--}10^{52} \text{ erg s}^{-1}$ , this would require an extremely long-lived phase of continuous accretion to explain hyper-energetic GRBs.

Jets powered by MHD processes, on the other hand, can more easily accommodate large energy releases. To begin with, by extracting energy directly from the rotation of the black hole, the Blandford-Znajek process begins with a significantly larger energy reservoir:  $E_{\text{rot}} \approx 8 \times 10^{53} \text{ erg}$  for a rapidly spinning ( $a = 0.9$ ) black hole of mass  $3 M_{\odot}$  (Eqn. 1). Both analytic models (e.g., Lee et al. 2000) and numerical simulations (e.g., McKinney 2005) suggest that as much as 5–10% of this rotational energy can be made available to power a collimated outflow, easily within the requirements of hyper-energetic events. We caution, however, that the jet efficiency is a strong function of the black hole spin, and slowly spinning black holes have significantly lower efficiencies.

Setting energetics considerations aside for the moment, there still remains the question of whether either the  $\nu\bar{\nu}$  annihilation mechanism or MHD processes are capable of producing jets with the Lorentz factors and degree of collimation inferred from GRB observations. Here our sample of *Fermi*-LAT events, with their extreme initial Lorentz factors, offers a distinct advantage over previous GRB studies. Combining the Lorentz-factor limits for the most relativistic GRBs with the inferred jet opening angles from our broadband afterglow models can provide critical diagnostics of the jet acceleration mechanism.

The results from MHD simulations of jet acceleration appear to depend sensitively on the nature of the medium into which the jet propagates. Tchekhovskoy et al. (2009b) have recently conducted fully relativistic MHD simulations in which a mildly magnetized (magnetization parameter  $\sigma \lesssim 1$ ) jet is initially confined by the pressure of a stellar envelope out to some radius, and then allowed to propagate freely in all directions (see also Komissarov et al. 2009). Both authors find that the jet accelerates rapidly in this transition region (“rarefaction” acceleration) and can reach Lorentz factors  $\Gamma_0$  of a few hundred, yet still remains highly collimated ( $\theta \lesssim 5^\circ$ ) even after leaving the region of confinement. Furthermore, the deconfinement radii required to produce these outflows agree well with the expected value for Wolf-Rayet stars ( $r \approx 10^9\text{--}10^{11} \text{ cm}$ ).

Based on both results from these simulations and analytical arguments, Tchekhovskoy et al. (2009b) demonstrate a relationship between the initial Lorentz factor and jet opening angle:  $\Gamma_0 \theta \approx 10\text{--}30$ . This result is able to nicely reproduce the typical observed Lorentz factors (Lithwick & Sari 2001) and opening angles (Zeh et al. 2006) of previous GRB samples. However, for the events studied here, both GRB 090902B ( $\Gamma_0 \theta \approx 70$ ) and GRB 090926A ( $\Gamma_0 \theta \approx 90$ ) appear inconsistent with this result.

One way to circumvent this requirement is if the jets have extremely high magnetization parameters ( $\sigma \gg 1$ ; Tchekhovskoy et al. 2009a), as such Poynting flux-




FIG. 10.— Two-dimensional relativistic energy release ( $E_{\text{rel}} \approx E_{\gamma} + E_{\text{KE}}$ ) from GRBs. Cosmologically distant ( $z \gtrsim 0.5$ ) events from the pre-*Swift* era are shown in red. The logarithmic mean for these events,  $\langle E_{\text{rel}} \rangle = 2 \times 10^{51}$  erg, is indicated by the solid black line. Shaded regions correspond to  $1\sigma$ ,  $2\sigma$ , and  $3\sigma$  errors on this mean value. The three most nearby events (GRBs 980425, 031203, and 060218) are plotted in green and are underluminous by several orders of magnitude. The four LAT events from this work are plotted in blue; all but GRB 090328 fall at the high end of the pre-*Swift* distribution (note that we have not plotted horizontal errors bars for GRB 090926A due to the large uncertainty in  $E_{\text{KE}}$ ). Instead, they are more consistent with some of the brightest events from the *Swift* era (black squares). The total relativistic energy release from GRB 090926A appears to exceed  $10^{52}$  erg. Such hyper-energetic events pose a severe challenge to the magnetar models, where the total energy release cannot exceed  $3 \times 10^{52}$  erg (dashed black line). References — Panaitescu & Kumar (2002): GRBs 990123, 990510, 991208, 991216, 000301C, 010222; Yost et al. (2003): GRBs 970508, 980703, 000926; Berger et al. (2004): GRBs 970508, 980703; Chevalier et al. (2004): GRB 020405; Berger et al. (2001): GRB 000418; Li & Chevalier (1999): GRB 980425; Soderberg et al. (2004): GRB 031203; Soderberg et al. (2006): GRB 060218; Frail et al. (2006): GRB 050904; Chandra et al. (2008): GRB 070125; Cenko et al. (2010): GRBs 050820A, 060418, 080319B.

dominated jets can be accelerated to extreme Lorentz factors over relatively large angles. However, it is unclear how such outflows can convert sufficient electromagnetic energy to accelerate electrons and produce the observed prompt gamma-ray emission. Other particle-acceleration mechanisms besides MHD shocks may be required in this case (e.g., Beloborodov 2009). Alternatively, the gamma-ray emission may be patchy (e.g., Kumar & Piran 2000) or the jet may be structured (see, e.g., Granot 2007 and references therein), so that we are measuring only the extrema of  $\Gamma_0$  and not the true bulk of the relativistic flow carrying most of the energy.

#### 4.3. Comparison with Other Work

In this section, we attempt to place this work in context, both by comparing our results with those of other authors who have studied these same events, and by highlighting additional differences between our LAT sample and GRBs detected by satellites at lower energies.

McBreen et al. (2010) present optical and NIR observations of three events from this work (GRBs 090323, 090328, and 090902B), taken primarily with the GROND instrument (Greiner et al. 2008). In the case of GRB 090323, these authors find an optical spectral in-

dex of  $\alpha_O = 1.90 \pm 0.01$ , consistent with the value derived here, and our measurements of the host-galaxy flux agree nicely. We derive a slightly steeper optical spectral index, but our results are consistent at the  $2.5\sigma$  level. Most importantly, McBreen et al. (2010) infer from the steep optical decay that a jet break occurred *before* the first optical observations began ( $t_j \lesssim 1$  day), resulting in a narrow beaming angle ( $\theta \lesssim 2^\circ$ ) and a correspondingly small collimation-corrected prompt energy release [ $E_{\gamma} \lesssim 3(1) \times 10^{51}$  erg for a constant-density (wind-like) circumburst medium]. We consider this possibility unlikely, however, as it is difficult to explain both the flat radio light curve and the more slowly fading X-ray afterglow ( $\alpha_X \approx 1.5$ ) through post jet-break evolution (see also § 4.4).

Our results also differ from those of McBreen et al. (2010) regarding the jet break time of GRB 090328. These authors argue that the steep optical decay index ( $\alpha_o \approx 2.3$ ) derived at early times requires a jet break before the commencement of observations ( $t_j \lesssim 1.5$  days). The data presented in our work are not sufficient to uniquely determine the optical temporal decay index (particularly given the possible contribution from an underlying host galaxy). However, it is again difficult to



reconcile the shallower X-ray decay ( $\alpha_X \approx 1.65$ ) and in particular the rising radio light curve at this time with models of post-jet break evolution (see also § 4.4).

We find very similar results to those of McBreen et al. (2010) for the temporal and spectral indices, inferred circumburst medium, and jet break time for GRB 090902B. In particular, the late-time ( $t \approx 23$  days) optical observation with the VLT provides strong confirmation that the radio decline at  $t \gtrsim 6$  days is due to a jet break. However, our results diverge when translating the jet break time into an opening angle. Because of the low density we derive for the circumburst medium of GRB 090902B, a jet break time of  $t_j \approx 6$  days corresponds to a relatively narrow opening angle. The resulting constraints on  $E_\gamma$  are roughly a factor of 4 less (note that part of this difference is also due to the wider bandpass McBreen et al. 2010 use to calculate  $E_{\gamma, \text{iso}}$ ).

Pandey et al. (2010) also present optical and X-ray observations of GRB 090902B, most of which were included in our broadband modeling here. It is not surprising, then, that we derive similar values for the optical temporal and spectral indices (the same also holds for the X-ray bandpass). The afterglow model derived by these authors is broadly similar to ours, but we favor a somewhat steeper electron index ( $p = 2.2$  vs. 1.8). Most importantly, Pandey et al. (2010) derive a limit on the opening angle of  $\theta > 6^\circ$  (based on a lower limit to the jet break time of  $t_j > 6$  days). Though our derived jet break time is consistent with the results of these authors, our inferred opening angle is a factor of two smaller due to the lower value we have used for the circumburst density. As a result, the limits we derive for the prompt gamma-ray energy release are a factor of a few less.

Kumar & Duran (2009a) (see also Kumar & Duran 2009b) have also presented an analysis of the broadband afterglow of GRB 090902B. Interestingly, these authors suggest that the delayed high-energy ( $E \gtrsim 100$  MeV) component observed from several LAT events is in fact due to the same source as the late-time afterglow emission: synchrotron radiation from accelerated electrons in the circumburst medium shock-heated by the outgoing blast wave (i.e., external shock emission; see also Ghisellini et al. 2010).

The afterglow parameters derived by these authors for GRB 090902B differ somewhat from ours, largely due to the fact that they assume the X-ray bandpass at  $t \approx 1$  day falls below the synchrotron cooling frequency (i.e.,  $\nu_X < \nu_c$ ). While the observed X-ray spectral and temporal indices at  $t \approx 1$  day are consistent with this picture, we find that a better fit can be achieved for both the X-ray and optical data if  $\nu_X > \nu_c$  at  $t \approx 1$  day (see also Pandey et al. 2010).

Using the best-fit late-time afterglow parameters we have derived for GRB 090902B (Table 6), we have attempted to calculate the magnitude of the afterglow (i.e., external shock) flux at  $\nu = 100$  MeV and  $t = 50$  s. We find  $f_\nu \approx 30$  nJy, roughly a factor of 7 below the observed value (Abdo et al. 2009a). The discrepancy between our results and those of Kumar & Duran (2009a) is largely due to the lower kinetic energy we have inferred for the blast wave ( $E_{\text{KE, iso}} \approx 7 \times 10^{53}$ , a factor of 18 lower than that used by Kumar & Duran 2009a). This would suggest that the delayed high-energy component is not due to afterglow emission. We caution, however, that our




FIG. 11.— Circumburst density and isotropic afterglow energy release from cosmological ( $z > 0.5$ ) GRBs. Constant-density models are shown as squares, while wind-like models are plotted as circles. The GRBs detected by the *Fermi*-LAT (red) clearly have preferentially larger isotropic energy releases and smaller circumburst densities than the rest of the sample. See the caption of Figure 10 for references.

flux calculation does not incorporate the reduced cross-section for inverse Compton emission due to the Klein-Nishina effect at very high energies, and this will affect our flux calculation to some extent.

A further interesting claim from the work of Kumar & Duran (2009a) is that the inferred value of the magnetic field for GRB 090902B (along with several other LAT-detected events) is consistent with shock compression of a modest circumstellar field ( $B \gtrsim 30 \mu\text{G}$ ). In other words, no dynamo process is necessary to generate the magnetic field strengths needed to produce the observed synchrotron afterglow emission. For a constant-density circumburst medium, the preshock magnetic field is given by

$$B = (2\pi m_p \epsilon_B n)^{1/2} c. \quad (2)$$

We also find that the product  $\epsilon_B \times n$  (and hence the derived  $B$  field) is smaller for LAT events than for other GRBs detected at lower energies, primarily due to the lower inferred circumburst densities (see below). However, using our best-fit parameters for GRB 090902B, we find  $B \approx 100 \mu\text{G}$ , broadly consistent with the results of Piran & Nakar (2010) and Li (2010). While this value is smaller than the preshock magnetic fields previously inferred for GRBs ( $B \approx 10$  mG for  $\epsilon_B = 0.01$ ,  $n = 1 \text{ cm}^{-3}$ ), it suggests that at least some magnetic field amplification may still be required.

Finally, we return to the issue of the relatively low circumburst densities derived in § 3. Given the strong evidence for a connection between long-duration GRBs and broad-lined SNe Ib/c, it would be natural to expect the massive-star progenitors of GRBs to explode in the dense molecular cloud environments where we observe stars forming in our own Galaxy.

Past modeling of broadband GRB afterglows, however, has not always revealed this to be the case. Of the past GRBs with sufficient radio observations to estimate the circumburst density, the derived values span a large range from  $1.9 \times 10^{-3} \text{ cm}^{-3}$  (GRB 990123; Panaitescu & Kumar 2002) to  $680 \text{ cm}^{-3}$  (GRB 050904; Frail et al. 2006). We plot in Figure 11 the derived cir-

cumburst densities for these previous events (either  $n$  or  $A_*$ , depending on the circumburst density profile) as a function of isotropic afterglow kinetic energy, along with three events from our LAT sample (we have not included GRB 090926A as lack of radio coverage makes density estimates highly degenerate). It is clear that the events detected by the LAT have on average larger isotropic energies and smaller densities than the previous sample.

The larger isotropic kinetic energy is not hard to understand, as the LAT is less sensitive than, for example, the Burst Alert Telescope (BAT; Barthelmy et al. 2005) onboard *Swift* and should therefore select brighter events (in terms of the high-energy fluence). Unless  $E_{\gamma,\text{iso}}$  and  $E_{\text{KE,iso}}$  were anticorrelated, we would expect LAT events to have larger isotropic blast-wave energies as well. But the lower densities require an additional explanation. We speculate on possible reasons for this difference in § 5.

The large ratio of  $E_{\text{KE,iso}}/n$  (or, alternatively  $E_{\text{KE,iso}}/A_*$ ) we have derived for these events has two important implications. First, the opening angle derived for a given jet break time scales inversely as  $E_{\text{KE,iso}}$  and proportionally to  $n/A_*$  (§ 4.4). Thus, for a given jet break time, the large ratios we have inferred for LAT events imply smaller opening angles than were one to simply use a canonical value (e.g.,  $n = 0.1 \text{ cm}^{-3}$  from Frail et al. 2001;  $n = 10 \text{ cm}^{-3}$  from Bloom et al. 2003). In this manner, LAT events can have relatively late jet breaks but still be narrowly collimated ( $\theta \lesssim 10^\circ$ ).

Secondly, a large  $E_{\text{KE,iso}}/n$  ratio will act to delay the deceleration time of the outflow. Assuming the “thin shell” case, the outgoing relativistic blast wave will decelerate when (Sari & Piran 1999; Molinari et al. 2007)

$$t_{\text{dec}}(\text{ISM}) = \left( \frac{3E_{\text{KE,iso}}(1+z)^3}{32\pi m_p c^5 n \Gamma_0^8} \right)^{1/3}, \quad (3)$$

$$t_{\text{dec}}(\text{Wind}) = \frac{E_{\text{KE,iso}}(1+z)}{8\pi c^3 A \Gamma_0^4}. \quad (4)$$

(Note that we have corrected Eqn. 2 from Molinari et al. 2007 to remove the erroneous factor of  $m_p$ .) To some extent, the large ratio of  $E_{\text{KE,iso}}/n$  will offset the effect of the larger initial Lorentz factors for *Fermi*-LAT events. Using the parameters we have derived from the LAT events in our sample, we find  $t_{\text{dec}} \approx 2 \text{ s}$  for GRB 090323,  $t_{\text{dec}} \approx 3 \text{ s}$  for GRB 090328, and  $t_{\text{dec}} \approx 33 \text{ s}$  for GRB 090902B. If the late-time GeV emission is indeed due to external shock emission, the delay between the MeV and GeV photons should correspond roughly to the blast-wave deceleration time. This may be problematic for GRB 090902B, for which the observed delay ( $\sim 4 \text{ s}$ ) is significantly less than our inferred deceleration time. We caution, however, that the large systematic uncertainties in  $E_{\text{KE,iso}}$ ,  $n$ , and particularly  $\Gamma_0$  (§ 4.4) may introduce a significant error into calculations of the deceleration time.

#### 4.4. Limitations and Future Work

To better understand our results in § 4.1, § 4.2, and § 4.3 on GRB energetics and circumburst density, it is important to consider the robustness of our modeling. The most crucial factor in determining the true energy release is the collimation correction, which we therefore examine in more detail here.

For a wind-like circumburst medium, the jet half-opening angle  $\theta$  can be written as (e.g., Chevalier & Li 2000)

$$\theta(\text{Wind}) = 0.10 \left( \frac{1+z}{2} \right)^{-1/4} \left( \frac{E_{\text{KE,iso}}}{10^{52} \text{ erg}} \right)^{-1/4} \left( \frac{A_*}{1.0 \text{ cm}^{-1}} \right)^{1/4} \left( \frac{t_j}{1 \text{ d}} \right)^{1/4} [\text{rad}]. \quad (5)$$

This result assumes only that the shock is ultrarelativistic and undergoes a self-similar evolution (Blandford & McKee 1976), and is therefore relatively robust.

Only in the case of GRB 090926A (and also for GRB 090902B if we include the late-time VLT observation from McBreen et al. 2010) do we find clear evidence for a jet break in multiple bandpasses. For GRB 090323, the lack of a steepening in the X-ray and optical light curves apparently limits  $t_j \gtrsim 10 \text{ days}$ . If we simply ignore our modeling results and assume that the shock energy is comparable to the prompt gamma-ray energy ( $E_{\text{KE,iso}} \approx 4 \times 10^{54} \text{ erg}$ ) and that the progenitor wind speed and mass-loss rate are comparable to those observed from Galactic Wolf-Rayet stars ( $A_* \approx 1 \text{ g cm}^{-1}$ ), then we find a lower limit on the opening angle of  $\theta(\text{Wind}) \gtrsim 3^\circ$ . The resulting limit on the prompt energy release is therefore  $E_\gamma \gtrsim 5 \times 10^{51} \text{ erg}$ , similar to what we derived from our broadband models.

If we were instead to assume the GRB exploded in a constant-density medium, the opening angle then becomes (e.g., Sari et al. 1999)

$$\theta(\text{ISM}) = 0.12 \left( \frac{1+z}{2} \right)^{-3/8} \left( \frac{E_{\text{KE,iso}}}{10^{52} \text{ erg}} \right)^{-1/8} \left( \frac{n}{1 \text{ cm}^{-3}} \right)^{1/8} \left( \frac{t_j}{1 \text{ d}} \right)^{3/8} [\text{rad}]. \quad (6)$$

Under similar assumptions as above, the limit on the opening angle would be  $\theta(\text{ISM}) \gtrsim 6^\circ$ , corresponding to a prompt energy release of  $E_\gamma \gtrsim 2 \times 10^{52} \text{ erg}$ . Clearly a constant-density circumburst medium only makes the energy requirements more strict.

An alternative possibility for GRB 090323 (McBreen et al. 2010; also suggested by Schady et al. 2007 to explain the afterglow of GRB 061007) is a jet break before the beginning X-ray and optical observations ( $t \lesssim 1 \text{ day}$  for GRB 090323). Much like GRB 061007, however, the implied post-break decay index in the X-ray ( $\alpha_X \approx 1.5$ ) is too shallow to accommodate a typical electron spectral index  $p$  (for  $\nu > \nu_m$ ,  $f_\nu \propto t^{-p}$  post jet-break; Sari et al. 1999). The observed behavior would therefore require late-time energy input to flatten the X-ray and optical decay. Furthermore, the relatively flat radio light curve for  $t \lesssim 20 \text{ days}$  would require that the synchrotron self-absorption frequency fell above the radio bandpass during this phase; otherwise the radio bandpass would be required to decay, either as  $t^{-1/3}$  ( $\nu < \nu_m$ ) or  $t^{-p}$  ( $\nu > \nu_m$ )<sup>30</sup>. We consider

<sup>30</sup> We note that the above results are independent of circumburst medium, as the expansion of the outflow is predominantly lateral after the jet break (Chevalier & Li 2000).

this somewhat contrived picture unlikely in the case of GRB 090323, particularly given the reasonable quality of our fits using only standard afterglow theory.

A similar analysis of GRB 090328 suggests that the jet-break time cannot occur any earlier than  $t \approx 4$  days, based again on the relatively steady decay seen in the X-ray light curve and, in particular, on the rising radio emission at early times. The corresponding limits on the opening angles are therefore  $\theta(\text{Wind}) \gtrsim 8^\circ$ ,  $\theta(\text{ISM}) \gtrsim 7^\circ$ . Both values result in a prompt energy release  $E_\gamma \gtrsim 10^{51}$  erg, broadly consistent with our modeling results.

In the end, all of these estimates of energy, density, and jet geometry rest on the relativistic synchrotron model. While the standard afterglow model has undergone continuous improvements and it has been well tested (Meszaros & Rees 1997; Sari et al. 1998, 1999; Wijers & Galama 1999; Chevalier & Li 2000; Panaitescu & Kumar 2000; Yost et al. 2003; Willingale et al. 2007), it still must make several simplifying assumptions about the shock dynamics, magnetic field generation, particle acceleration, energy injection, and the circumburst density structure. Of particular concern are the recent results of relativistic simulations by Zhang & MacFadyen (2009), which suggest that viewing angle can dramatically alter the observed afterglow emission at early times.

If we are to verify this class of hyper-energetic bursts, we need independent estimates of the relativistic energy content. Waxman et al. (1998) first suggested that late-time (radio) calorimetry could be used for this purpose in order to sidestep early-time complications such as the outflow geometry, the density structure, and ongoing activity from the central engine. Recent relativistic hydrodynamic simulations show that the earlier analytic models which used spherical geometry and Sedov-Taylor dynamics to explain the late-time behavior were approximately correct and produce a robust estimate of the total kinetic energy (Zhang & MacFadyen 2009). In the past, the method has been limited to only a small number of radio-bright events (e.g., van der Horst et al. 2008 and references therein). But it is somewhat encouraging that both early- and late-time estimates for the most energetic pre-*Fermi* event (GRB 970508;  $E_{\text{KE}} \approx 10^{52}$  erg) are in close agreement (Yost et al. 2003; Berger et al. 2004). With the advent of radio facilities having greatly increased sensitivity such as LOFAR (van Haarlem 2005) and the EVLA (Perley et al. 2009c), it should soon be possible to verify candidate hyper-energetic events by searching for long-lived ( $t \gtrsim 100$  days) radio afterglows.

Finally, we add a further note of caution that the derivation of the initial Lorentz factor  $\Gamma_0$  is also subject to significant systematic uncertainties (e.g., Bošnjak et al. 2009). In order to calculate the optical depth to pair production, it is necessary to determine the effective blast-wave radius as well as the instantaneous spectral parameters. The former is typically inferred from the variability time scale, while the latter is usually averaged over the entire duration of the prompt emission, and it is not always clear how large an effect this will have on the calculation of  $\Gamma_0$ . The calculations are further complicated if the spectrum is not a pure power law, as would be the case for thermal emission arising from the jet photosphere (Ryde et al. 2010). Here we

have attempted to calculate the initial Lorentz factors in a consistent manner, and it is not unreasonable to believe that the relative ordering of  $\Gamma_0$  is robust. However, small changes in  $\Gamma_0$  can have a large effect on some of the parameters derived here (e.g.,  $t_{\text{dec}}$ ; Eqns. 3 and 4). Future early afterglow observations of *Fermi*-LAT GRBs to directly measure the deceleration time and hence constrain  $\Gamma_0$  (e.g., Molinari et al. 2007) could provide valuable insight in this area.

## 5. CONCLUSIONS

We have undertaken extensive broadband continuum (radio, optical, and X-ray) and spectroscopic observations of four long-duration GRBs (GRBs 090323, 090328, 090902B, and 090926A) detected by the LAT instrument on the *Fermi* satellite at GeV energies. This work was motivated by the realization that *Fermi* is especially sensitive to GRBs with large isotropic energy release, and hence provides an interesting sample of events to test GRB central-engine models and their relativistic outflows. Our afterglow models constrain the jet break times and the density of the circumburst medium, from which we derive the collimation angle and hence beaming-corrected energy release for each event. We find three GRBs with a total relativistic content about an order of magnitude in excess of the canonical  $10^{51}$  erg, with GRB 090926A almost certainly in excess of  $10^{52}$  erg. This analysis provides support for our earlier claim of a class of hyper-energetic GRBs. The discovery of more GRBs with total energy release  $\gtrsim 10^{52}$  erg is troubling for central engines in which the energy to drive the jet is derived either from a rotating magnetar or collapsars powered by neutrino annihilation. For this reason, we are led to believe that, at least for hyper-energetic GRBs, the massive star progenitor collapses directly to a black hole and the rotational energy of this system is extracted via the Blandford-Znajek process.

Although we find relatively narrow opening angles for all four events ( $\theta \lesssim 10^\circ$ ), the extreme initial Lorentz factors inferred for these LAT events imply that the product  $\theta\Gamma_0$  can be a factor of 5–10 larger than estimates of previous GRBs detected at MeV energies. These values are inconsistent with recent simulations of low-magnetization MHD jets, suggesting that the outflow may be at least initially Poynting-flux dominated. If this is indeed the case, it is unclear how the initial kinetic energy of the outflow is converted to prompt gamma-ray emission.

Interestingly, for the three events having sufficient radio coverage to derive a circumburst density, we find anomalously large values of  $E_{\text{KE,iso}}/n$  (or, for a wind-like medium,  $E_{\text{KE,iso}}/A_*$ ). While the large  $E_{\text{KE,iso}}$  values are simple to understand, the low circumburst densities require a more complex explanation.

One possibility is that the progenitor stars of LAT GRBs are somehow different from the progenitors of most previous GRBs detected at MeV or keV energies. It is currently thought that GRB progenitors are distinguished from the progenitors of ordinary SNe Ib/c by their low metallicities (e.g., MacFadyen & Woosley 1999; Woosley & Heger 2006; Modjaz et al. 2008): the lower mass-loss rates allow the progenitors of GRBs to keep more angular momentum. The increased rotation evacuates a cavity through which a relativistic jet can propagate. If LAT events have larger initial Lorentz factors,

it may be that they come from lower metallicity progenitors with minimal pre-explosion mass loss. Observations of the host galaxies of these events, both through absorption and emission spectroscopy, may help shed light on this matter.

It is also possible that the low density preference is the result of other, more subtle, selection effects. In particular, if the GeV emission arises in the external shock, the *Fermi*-LAT could be biased towards events in low-density environments. If the circumburst density is too high, the blastwave will decelerate at small radii (Equations 3 and 4), where the outflow may be opaque to GeV photons. More observations of LAT GRBs, particularly at very early times, would help to investigate this hypothesis.

We end by emphasizing the importance of afterglow observations of high- $E_{\gamma, \text{iso}}$  events in the *Fermi* era to provide further confirmation of this picture. Such GRBs are either highly collimated outflows ( $\theta \lesssim 2^\circ$ ) with a typical energy release, or truly hyper-energetic events; both represent extreme tests of jet collimation and central-engine models, respectively. Current efforts suffer from delays in LAT localizations and limited ground-based afterglow follow-up efforts. The latter can be improved by focusing rare follow-up resources on *Fermi*-LAT GRBs; as Nysewander et al. (2009) and McBreen et al. (2010) have shown, these events have brighter X-ray and optical afterglows on average, and are therefore accessible even for moderate-aperture optical facilities. Targeting these bright afterglows will make it easier to measure the jet breaks, which have proven almost impossible to obtain in the *Swift* era. Finally, we note that one testable consequence of hyper-energetic GRBs is long-lived afterglow emission ( $\gtrsim 1$  yr). If the shock microphysics and the circumburst density do not undergo a drastic evolution, it should be possible to detect these afterglows with the upcoming generation of radio facilities and carry out

calorimetry measurements.

S.B.C. and A.V.F. wish to acknowledge generous support from Gary and Cynthia Bengier, the Richard and Rhoda Goldman Fund, National Aeronautics and Space Administration (NASA)/*Swift* grants NNX09AL08G and NNX10AI21G, and National Science Foundation (NSF) grants AST-0607485 and AST-0908886. B.E.C. gratefully acknowledges support from an NSF Astronomy & Astrophysics Postdoctoral Fellowship (AST-0802333). N.R.B. is supported through the Einstein Fellowship Program (NASA Cooperative Agreement NNG06DO90A). J. S. B. and his group were partially supported by NASA/*Swift* Guest Investigator grant NNX09AQ66G and a grant from DOE SciDAC.

P60 operations are funded in part by NASA through the *Swift* Guest Investigator Program (grant number NNG06GH61G). Based in part on observations obtained at the Gemini Observatory (Programs GS-2009A-Q-23, GS-2009B-Q-5, GN-2009A-Q-26, and GN-2009B-Q-28), which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (US), the Particle Physics and Astronomy Research Council (UK), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), CNPq (Brazil) and CONICET (Argentina). We wish to thank the entire staff at Gemini for assistance with these observations. SMARTS is supported by NSF grant AST-0707627.

*Facilities:* Fermi (LAT, GBM), VLA, PO:1.5m, Gemini:South (GMOS), Gemini:North (GMOS), CTIO:2MASS (ANDICAM), Swift (XRT, UVOT)

## REFERENCES

- Abazajian, K. N., et al. 2009, *ApJS*, 182, 543  
 Abdo, A. A., et al. 2009a, *ApJ*, 706, L138  
 —. 2009b, *Science*, 323, 1688  
 Amati, L. 2006, *MNRAS*, 372, 233  
 Asano, K., Guiriec, S., & Mészáros, P. 2009, *ApJ*, 705, L191  
 Atwood, W. B., et al. 2009, *ApJ*, 697, 1071  
 Baldwin, J. A., & Stone, R. P. S. 1984, *MNRAS*, 206, 241  
 Band, D., et al. 1993, *ApJ*, 413, 281  
 Barthelmy, S. D., et al. 2005, *Space Science Reviews*, 120, 143  
 Beloborodov, A. M. 2009, eprint arXiv, (astro-ph/0907.0732)  
 Berger, E., Kulkarni, S. R., & Frail, D. A. 2003a, *ApJ*, 590, 379  
 —. 2004, *ApJ*, 612, 966  
 Berger, E., Soderberg, A. M., Frail, D. A., & Kulkarni, S. R. 2003b, *ApJ*, 587, L5  
 Berger, E., et al. 2001, *ApJ*, 556, 556  
 —. 2005, *ApJ*, 634, 501  
 Birkel, R., Aloy, M. A., Janka, H.-T., & Müller, E. 2007, *A&A*, 463, 51  
 Bissaldi, E. 2009, GRB Coordinates Network, 9933, 1  
 Bissaldi, E., Briggs, M. S., Piron, F., Takahashi, H., & Uehara, T. 2009, GRB Coordinates Network, 9972, 1  
 Bissaldi, E., & Connaughton, V. 2009, GRB Coordinates Network, 9866, 1  
 Blandford, R. D., & McKee, C. F. 1976, *Physics of Fluids*, 19, 1130  
 Blandford, R. D., & Znajek, R. L. 1977, *MNRAS*, 179, 433  
 Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, *ApJ*, 594, 674  
 Bloom, J. S., Frail, D. A., & Sari, R. 2001, *The Astronomical Journal*, 121, 2879  
 Bošnjak, Ž., Daigne, F., & Dubus, G. 2009, *A&A*, 498, 677  
 Burenin, R., et al. 2009, GRB Coordinates Network, 9037, 1  
 Burrows, D. N., et al. 2005a, *Science*, 309, 1833  
 —. 2005b, *Space Science Reviews*, 120, 165  
 Butler, N. R., Bloom, J. S., & Poznanski, D. 2010, *ApJ*, 711, 495  
 Butler, N. R., & Kocevski, D. 2007, *ApJ*, 663, 407  
 Butler, N. R., Kocevski, D., Bloom, J. S., & Curtis, J. L. 2007, *ApJ*, 671, 656  
 Campana, S., et al. 2006, *Nature*, 442, 1008  
 Cavallo, G., & Rees, M. J. 1978, *MNRAS*, 183, 359  
 Cenko, S. B., Bloom, J. S., Morgan, A. N., & Perley, D. A. 2009, GRB Coordinates Network, 9053, 1  
 Cenko, S. B., & Perley, D. A. 2009, GRB Coordinates Network, 9027, 1  
 Cenko, S. B., et al. 2006, *PASP*, 118, 1396  
 —. 2008, *ApJ*, 677, 441  
 —. 2010, *ApJ*, 711, 641  
 Chakrabarti, S. K., et al. 2009, GRB Coordinates Network, 10009, 1  
 Chandra, P., & Frail, D. A. 2009, GRB Coordinates Network, 9889, 1  
 Chandra, P., et al. 2008, *ApJ*, 683, 924  
 —. 2010, *ApJ*, 712, L31  
 Chandra, P., et al. 2010, in preparation  
 Chevalier, R. A., & Li, Z.-Y. 1999, *The Astrophysical Journal*, 520, L29  
 —. 2000, *ApJ*, 536, 195  
 Chevalier, R. A., Li, Z.-Y., & Fransson, C. 2004, *ApJ*, 606, 369  
 Chornock, R., Perley, D. A., Cenko, S. B., & Bloom, J. S. 2009, GRB Coordinates Network, 9028, 1  
 Cobb, B. E., Bailyn, C. D., van Dokkum, P. G., Buxton, M. M., & Bloom, J. S. 2004, *ApJ*, 608, L93

- Cobb, B. E., Baily, C. D., van Dokkum, P. G., & Natarajan, P. 2006, *ApJ*, 645, L113
- Cucchiara, A., Fox, D. B., Tanvir, N., & Berger, E. 2009, GRB Coordinates Network, 9873, 1
- Curran, P. A., Starling, R. L. C., van der Horst, A. J., & Wijers, R. A. M. J. 2009, *MNRAS*, 395, 580
- Cutini, S., Vasileiou, V., & Chiang, J. 2009, GRB Coordinates Network, 9077, 1
- Daigne, F., & Mochkovitch, R. 1998, *MNRAS*, 296, 275
- de Palma, F., Bregeon, J., & Tajima, H. 2009, GRB Coordinates Network, 9867, 1
- de Ugarte Postigo, A., Xu, D., Malesani, D., Hjorth, J., Fynbo, J. P. U., Jakobsson, P., & Adamo, A. 2009, GRB Coordinates Network, 9051, 1
- Djorgovski, S. G., et al. 1997, *Nature*, 387, 876
- Evans, P. A. 2009, GRB Coordinates Network, 9871, 1
- Falcone, A. D., et al. 2007, *ApJ*, 671, 1921
- Ferrero, P., et al. 2006, *A&A*, 457, 857
- Frail, D. A., Chandra, P., & Cenko, B. 2009, GRB Coordinates Network, 9060, 1
- Frail, D. A., Soderberg, A. M., Kulkarni, S. R., Berger, E., Yost, S., Fox, D. W., & Harrison, F. A. 2005, *ApJ*, 619, 994
- Frail, D. A., Waxman, E., & Kulkarni, S. R. 2000a, *ApJ*, 537, 191
- Frail, D. A., et al. 2000b, *ApJ*, 538, L129
- . 2001, *ApJ*, 562, L55
- . 2006, *ApJ*, 646, L99
- Freedman, D. L., & Waxman, E. 2001, *ApJ*, 547, 922
- Fukugita, M., Shimasaku, K., & Ichikawa, T. 1995, *PASP*, 107, 945
- Gal-Yam, A., et al. 2004, *ApJ*, 609, L59
- Galama, T. J., et al. 1998, *Nature*, 395, 670
- Gehrels, N., Ramirez-Ruiz, E., & Fox, D. B. 2009, *ARA&A*, 47, 567
- Gehrels, N., et al. 2004, *ApJ*, 611, 1005
- Ghisellini, G., Ghirlanda, G., Nava, L., & Celotti, A. 2010, *Monthly Notices of the Royal Astronomical Society*, 403, 926, (c) Journal compilation © 2010 RAS
- Giuliani, A., et al. 2008, *A&A*, 491, L25
- Golenetskii, S., et al. 2009a, GRB Coordinates Network, 9049, 1
- . 2009b, GRB Coordinates Network, 9959, 1
- . 2009c, GRB Coordinates Network, 9030, 1
- . 2009d, GRB Coordinates Network, 9050, 1
- González, M. M., Dingus, B. L., Kaneko, Y., Preece, R. D., Dermer, C. D., & Briggs, M. S. 2003, *Nature*, 424, 749
- Granot, J. 2007, *Triggering Relativistic Jets* (Eds. William H. Lee & Enrico Ramírez-Ruiz) *Revista Mexicana de Astronomía y Astrofísica (Serie de Conferencias)*, 27, 140
- Granot, J., & Sari, R. 2002, *ApJ*, 568, 820
- Greiner, J., et al. 2008, *PASP*, 120, 405
- . 2009, *A&A*, 498, 89
- Grupe, D., et al. 2007, *ApJ*, 662, 443
- . 2010, *ApJ*, 711, 1008
- Guetta, D., & Valle, M. D. 2007, *ApJ*, 657, L73
- Guidorzi, C., et al. 2009, GRB Coordinates Network, 9039, 1
- Haislip, J., et al. 2009a, GRB Coordinates Network, 9953, 1
- . 2009b, GRB Coordinates Network, 9937, 1
- . 2009c, GRB Coordinates Network, 9984, 1
- . 2009d, GRB Coordinates Network, 0003, 1
- . 2009e, GRB Coordinates Network, 9982, 1
- Harikae, S., Kotake, K., & Takiwaki, T. 2010, *ApJ*, 713, 304
- Harrison, F., Cenko, B., Frail, D. A., Chandra, P., & Kulkarni, S. 2009, GRB Coordinates Network, 9043, 1
- Hook, I. M., Jørgensen, I., Allington-Smith, J. R., Davies, R. L., Metcalfe, N., Murowinski, R. G., & Crampton, D. 2004, *PASP*, 116, 425
- Hurley, K., Briggs, M. S., Kippen, R. M., Kouveliotou, C., Meegan, C., Fishman, G., Cline, T., & Boer, M. 1999, *ApJS*, 120, 399
- Hurley, K., Goldsten, J., von Kienlin, A., Lichti, G., Rau, A., Connaughton, V., Briggs, M., & Meegan, C. 2009, GRB Coordinates Network, 9023, 1
- Hurley, K., et al. 1994, *Nature*, 372, 652
- Iwamoto, K., et al. 1998, *Nature*, 395, 672
- Jakobsson, P., et al. 2006, *A&A*, 447, 897
- Jordi, K., Grebel, E. K., & Ammon, K. 2006, *A&A*, 460, 339
- Kann, D. A., Klose, S., & Zeh, A. 2006, *The Astrophysical Journal*, 641, 993
- Kann, D. A., Laux, U., Ludwig, F., & Stecklum, S. 2009a, GRB Coordinates Network, 9063, 1
- Kann, D. A., Laux, U., & Stecklum, S. 2009b, GRB Coordinates Network, 9033, 1
- . 2009c, GRB Coordinates Network, 9041, 1
- Kann, D. A., et al. 2007, eprint arXiv, (astro-ph/0712.2186)
- Kennea, J. 2009, GRB Coordinates Network, 9045, 1
- Kennea, J., Evans, P., & Goad, M. 2009a, GRB Coordinates Network, 9024, 1
- . 2009b, GRB Coordinates Network, 9046, 1
- Kennea, J. A., & Stratta, G. 2009, GRB Coordinates Network, 9868, 1
- Kobayashi, S., Piran, T., & Sari, R. 1997, *ApJ*, 490, 92
- Kocevski, D., & Butler, N. 2008, *ApJ*, 680, 531
- Komissarov, S. S., Vlahakis, N., & Konigl, A. 2009, eprint arXiv, (astro-ph/0912.0845)
- Kulkarni, S. R., et al. 1998, *Nature*, 395, 663
- . 1999, *ApJ*, 522, L97
- Kumar, P., & Duran, R. B. 2009a, eprint arXiv, (astro-ph/0910.5726)
- . 2009b, *MNRAS*, 400, L75
- Kumar, P., & Piran, T. 2000, *ApJ*, 535, 152
- Landolt, A. U. 1992, *AJ*, 104, 340
- Lazzati, D., Rossi, E., Covino, S., Ghisellini, G., & Malesani, D. 2002, *A&A*, 396, L5
- Lee, H. K., Wijers, R. A. M. J., & Brown, G. E. 2000, *Physics Reports*, 325, 83
- Li, W., Jha, S., Filippenko, A. V., Bloom, J. S., Pooley, D., Foley, R. J., & Perley, D. A. 2006, *PASP*, 118, 37
- Li, Z. 2010, eprint arXiv, 1004, 791, 6 pages
- Li, Z.-Y., & Chevalier, R. A. 1999, *ApJ*, 526, 716
- Liang, E., Zhang, B., Virgili, F., & Dai, Z. G. 2007, *ApJ*, 662, 1111
- Liang, E.-W., Racusin, J. L., Zhang, B., Zhang, B.-B., & Burrows, D. N. 2008, *ApJ*, 675, 528
- Lithwick, Y., & Sari, R. 2001, *ApJ*, 555, 540
- MacFadyen, A. I., & Woosley, S. E. 1999, *ApJ*, 524, 262
- MacFadyen, A. I., Woosley, S. E., & Heger, A. 2001, *ApJ*, 550, 410
- Maeda, K., et al. 2007, *ApJ*, 658, L5
- Malesani, D., et al. 2004, *ApJ*, 609, L5
- . 2009, GRB Coordinates Network, 9942, 1
- Marshall, F. E., Oates, S. R., Rowlinson, A., Barthelmy, S. D., Burrows, D. N., Roming, P., & Gehrels, N. 2009, GRB Coordinates Network Report, 207, 1
- Matheson, T., Filippenko, A. V., Ho, L. C., Barth, A. J., & Leonard, D. C. 2000, *AJ*, 120, 1499
- Mazzali, P. A., et al. 2006, *Nature*, 442, 1018
- McBreen, S., et al. 2010, eprint arXiv, (astro-ph/1003.3885)
- McEnery, J., Cutini, S., Ohno, M., & Koering, E. 2009, GRB Coordinates Network, 9044, 1
- McKinney, J. C. 2005, *ApJ*, 630, L5
- Meegan, C., et al. 2009, *ApJ*, 702, 791
- Mészáros, P., & Rees, M. J. 1997, *ApJ*, 476, 232
- Mészáros, P., & Rees, M. J. 2000, *ApJ*, 530, 292
- Metzger, B. D., Thompson, T. A., & Quataert, E. 2007, *ApJ*, 659, 561
- Mirabal, N., Halpern, J. P., An, D., Thorstensen, J. R., & Terndrup, D. M. 2006, *ApJ*, 643, L99
- Modjaz, M., et al. 2006, *ApJ*, 645, L21
- . 2008, *AJ*, 135, 1136
- Moin, A., Frail, D. A., Tingay, S., & Macquart, J. 2009, GRB Coordinates Network, 10020, 1
- Molinari, E., et al. 2007, *A&A*, 469, L13
- Monet, D. G., et al. 2003, *AJ*, 125, 984
- Nakar, E., & Granot, J. 2007, *MNRAS*, 380, 1744
- Narayan, R., Piran, T., & Kumar, P. 2001, *ApJ*, 557, 949
- Noda, K., et al. 2009, GRB Coordinates Network, 9951, 1
- Nysewander, M., Fruchter, A. S., & Pe'er, A. 2009, *ApJ*, 701, 824
- Oates, S. R. 2009, GRB Coordinates Network, 9048, 1
- Ohno, M., Cutini, S., McEnery, J., Chiang, J., & Koering, E. 2009, GRB Coordinates Network, 9021
- Oke, J. B. 1990, *AJ*, 99, 1621
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Panaiteescu, A. 2005, *MNRAS*, 363, 1409
- . 2006, *MNRAS*, 367, L42
- . 2007, *MNRAS*, 380, 374



- Panaitescu, A., & Kumar, P. 2000, *ApJ*, 543, 66  
—, 2001a, *ApJ*, 560, L49  
—, 2001b, *ApJ*, 554, 667  
—, 2002, *ApJ*, 571, 779
- Pandey, S. B., et al. 2010, eprint arXiv, (astro-ph/1003.4250)
- Pei, Y. C. 1992, *The Astrophysical Journal*, 395, 130
- Perley, D. A. 2009, *GRB Coordinates Network*, 9042, 1
- Perley, D. A., Klein, C. R., Morgan, A. N., & Petigura, E. 2009a, *GRB Coordinates Network*, 9036, 1
- Perley, D. A., et al. 2008, *ApJ*, 688, 470  
—, 2009b, *ApJ*, 696, 1871
- Perley, R., et al. 2009c, *Proceedings of the IEEE* (ISSN: 0018-9219), 97, 1448
- Perna, R., Sari, R., & Frail, D. 2003, *ApJ*, 594, 379
- Pian, E., et al. 2006, *Nature*, 442, 1011
- Piran, T. 2005, *Reviews of Modern Physics*, 76, 1143
- Piran, T., & Nakar, E. 2010, eprint arXiv, (astro-ph/1003.5919)
- Piron, F., et al. 2009, in 2009 *Fermi* Symposium, ed. eConf Proceedings C0911022
- Popham, R., Woosley, S. E., & Fryer, C. 1999, *ApJ*, 518, 356
- Price, P. A., et al. 2002, *ApJ*, 572, L51
- Racusin, J. L., et al. 2009, *ApJ*, 698, 43
- Rau, A., Connaughton, V., & Briggs, M. 2009, *GRB Coordinates Network*, 9057, 1
- Razzaque, S., Dermer, C. D., & Finke, J. D. 2009, eprint arXiv, (astro-ph/0908.0513)
- Rees, M. J., & Meszaros, P. 1998, *ApJ*, 496, L1
- Reichart, D., et al. 2005, *Il Nuovo Cimento C*, 28, 767
- Rhoads, J. E. 1999, *ApJ*, 525, 737
- Roming, P. W. A., et al. 2005, *Space Science Reviews*, 120, 95
- Rosswog, S., Ramirez-Ruiz, E., & Davies, M. B. 2003, *MNRAS*, 345, 1077
- Ruffert, M., & Janka, H.-T. 1999, *A&A*, 344, 573
- Rumyantsev, V., & Pozanenko, A. 2009, *GRB Coordinates Network*, 9324, 1
- Ryde, F. 2004, *ApJ*, 614, 827
- Ryde, F., et al. 2010, *ApJ*, 709, L172
- Sari, R., & Esin, A. A. 2001, *ApJ*, 548, 787
- Sari, R., & Piran, T. 1999, *ApJ*, 517, L109
- Sari, R., Piran, T., & Halpern, J. P. 1999, *ApJ*, 519, L17
- Sari, R., Piran, T., & Narayan, R. 1998, *ApJ*, 497, L17
- Sazonov, S. Y., Lutovinov, A. A., & Sunyaev, R. A. 2004, *Nature*, 430, 646
- Schady, P., et al. 2007, *MNRAS*, 380, 1041
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Shen, R., Kumar, P., & Robinson, E. L. 2006, *MNRAS*, 371, 1441
- Skrutskie, M. F., et al. 2006, *AJ*, 131, 1163
- Soderberg, A. M., & Ramirez-Ruiz, E. 2003, *MNRAS*, 345, 854
- Soderberg, A. M., et al. 2004, *Nature*, 430, 648  
—, 2006, *Nature*, 442, 1014  
—, 2010, *Nature*, 463, 513
- Sollerman, J., et al. 2006, *A&A*, 454, 503
- Spergel, D. N., et al. 2007, *ApJS*, 170, 377
- Starling, R. L. C., van der Horst, A. J., Rol, E., Wijers, R. A. M. J., Kouveliotou, C., Wiersema, K., Curran, P. A., & Weltevrede, P. 2008, *ApJ*, 672, 433
- Starling, R. L. C., et al. 2009, *MNRAS*, 400, 90
- Stone, R. P. S., & Baldwin, J. A. 1983, *MNRAS*, 204, 347
- Stratta, G., D’Elia, V., & Perri, M. 2009, *GRB Coordinates Network*, 9876, 1
- Swenson, C. A., & Siegel, M. H. 2009, *GRB Coordinates Network*, 9869, 1
- Swenson, C. A., & Stratta, G. 2009, *GRB Coordinates Network*, 9877, 1
- Tam, P. H., Pun, C. S. J., Huang, Y. F., & Cheng, K. S. 2005, *New Astronomy*, 10, 535
- Tchekhovskoy, A., McKinney, J. C., & Narayan, R. 2009a, *ApJ*, 699, 1789
- Tchekhovskoy, A., Narayan, R., & McKinney, J. C. 2009b, eprint arXiv, (astro-ph/0909.0011)
- Terada, Y., et al. 2009, *GRB Coordinates Network*, 9897, 1
- Thompson, T. A., Chang, P., & Quataert, E. 2004, *ApJ*, 611, 380
- Thomsen, B., et al. 2004, *A&A*, 419, L21
- Uehara, T., Takahashi, H., & McEnery, J. 2009, *GRB Coordinates Network*, 9934, 1
- Udike, A., Klose, S., Clemens, C., & Greiner, J. 2009a, *GRB Coordinates Network*, 9054, 1
- Udike, A. C., Filgas, R., Kruehler, T., Greiner, J., & McBreen, S. 2009b, *GRB Coordinates Network*, 9026, 1
- Usov, V. V. 1992, *Nature*, 357, 472
- van der Horst, A. J. 2009, *GRB Coordinates Network*, 9047, 1
- van der Horst, A. J., Kamble, A. P., Wijers, R. A. M. J., & Kouveliotou, C. 2009, *GRB Coordinates Network*, 9883, 1
- van der Horst, A. J., & Xin, L. P. 2009, *GRB Coordinates Network*, 9035, 1
- van der Horst, A. J., et al. 2008, *A&A*, 480, 35
- van Haarlem, M. P. 2005, *EAS Publications Series*, 15, 431
- Vetere, L. 2009, *GRB Coordinates Network*, 9961, 1
- Vetere, L., Evans, P. A., & Goad, M. R. 2009, *GRB Coordinates Network*, 9936, 1
- Wade, R. A., & Horne, K. 1988, *ApJ*, 324, 411
- Wang, X. F., Xin, L. P., Zheng, W. K., Qiu, Y. L., Wei, J. Y., Deng, J. S., & Hu, J. Y. 2009, *GRB Coordinates Network*, 9034, 1
- Waxman, E., Kulkarni, S. R., & Frail, D. A. 1998, *ApJ*, 497, 288
- Wijers, R. A. M. J., & Galama, T. J. 1999, *ApJ*, 523, 177
- Willingale, R., et al. 2007, *ApJ*, 662, 1093
- Woosley, S. E. 1993, *ApJ*, 405, 273
- Woosley, S. E., & Bloom, J. S. 2006, *ARA&A*, 44, 507
- Woosley, S. E., & Heger, A. 2006, *ApJ*, 637, 914
- Yost, S. A., Harrison, F. A., Sari, R., & Frail, D. A. 2003, *ApJ*, 597, 459
- Zeh, A., Klose, S., & Kann, D. A. 2006, *ApJ*, 637, 889
- Zhang, W., & MacFadyen, A. 2009, *ApJ*, 698, 1261
- Zhang, W., Woosley, S. E., & MacFadyen, A. I. 2003, *ApJ*, 586, 356

TABLE 9  
OPTICAL/NIR OBSERVATIONS OF GRB 090323

Date <sup>a</sup> (UT)	Time Since Burst <sup>b</sup> (days)	Telescope/Instrument	Filter	Exposure Time (s)	Magnitude <sup>c</sup>	Reference <sup>d</sup>
2009 Mar 24.12	1.12	GROND	<i>g'</i>	480.0	21.64 ± 0.07	1
2009 Mar 24.12	1.12	GROND	<i>r'</i>	480.0	20.03 ± 0.03	1
2009 Mar 24.12	1.12	GROND	<i>i'</i>	480.0	19.64 ± 0.02	1
2009 Mar 24.12	1.12	GROND	<i>z'</i>	480.0	19.39 ± 0.02	1
2009 Mar 24.12	1.12	GROND	<i>J</i>	480.0	19.24 ± 0.02	1
2009 Mar 24.12	1.12	GROND	<i>H</i>	480.0	18.86 ± 0.02	1
2009 Mar 24.12	1.12	GROND	<i>K<sub>s</sub></i>	480.0	18.58 ± 0.03	1
2009 Mar 24.19	1.20	P60	<i>r'</i>	1800.0	20.50 ± 0.07	*
2009 Mar 24.20	1.21	P60	<i>i'</i>	1800.0	19.95 ± 0.04	*
2009 Mar 24.23	1.23	Gemini-S/GMOS	<i>i'</i>	90.0	20.04 ± 0.03	*
2009 Mar 24.32	1.32	P60	<i>r'</i>	900.0	20.46 ± 0.04	*
2009 Mar 24.33	1.33	P60	<i>i'</i>	900.0	20.21 ± 0.04	*
2009 Mar 24.34	1.35	P60	<i>r'</i>	900.0	20.56 ± 0.04	*
2009 Mar 24.36	1.36	P60	<i>i'</i>	900.0	20.20 ± 0.04	*
2009 Mar 24.42	1.42	P60	<i>r'</i>	900.0	20.65 ± 0.05	*
2009 Mar 24.43	1.43	P60	<i>i'</i>	900.0	20.36 ± 0.05	*
2009 Mar 24.44	1.44	P60	<i>r'</i>	900.0	20.60 ± 0.04	*
2009 Mar 24.45	1.46	P60	<i>i'</i>	900.0	20.34 ± 0.06	*
2009 Mar 24.65	1.65	Xinglong/TNT	<i>R</i>	6000.0	20.71	2
2009 Mar 24.88	1.88	TLS Tautenberg	<i>R</i>	4800.0	21.04 ± 0.04	3
2009 Mar 25.06	2.06	RTT150	<i>R</i>	3600.0	21.01 ± 0.04	4
2009 Mar 25.21	2.21	P60	<i>r'</i>	720.0	21.51 ± 0.10	*
2009 Mar 25.22	2.22	P60	<i>i'</i>	900.0	20.96 ± 0.09	*
2009 Mar 25.23	2.24	P60	<i>r'</i>	900.0	21.54 ± 0.11	*
2009 Mar 25.24	2.25	P60	<i>i'</i>	900.0	21.05 ± 0.09	*
2009 Mar 25.26	2.26	P60	<i>r'</i>	720.0	21.76 ± 0.09	*
2009 Mar 25.27	2.27	P60	<i>i'</i>	720.0	21.24 ± 0.08	*
2009 Mar 25.28	2.28	P60	<i>r'</i>	720.0	21.72 ± 0.08	*
2009 Mar 25.29	2.29	P60	<i>i'</i>	900.0	21.33 ± 0.08	*
2009 Mar 25.30	2.30	Nickel	<i>R</i>	3300.0	21.46 ± 0.2	5
2009 Mar 25.30	2.31	P60	<i>r'</i>	720.0	21.61 ± 0.07	*
2009 Mar 25.31	2.32	P60	<i>i'</i>	720.0	21.25 ± 0.07	*
2009 Mar 25.50	2.50	Faulkes South	<i>i'</i>	1800.0	21.3 ± 0.2	6
2009 Mar 25.53	2.53	Faulkes South	<i>R</i>	1800.0	21.7 ± 0.1	6
2009 Mar 26.23	3.25	P60	<i>r'</i>	3240.0	22.29 ± 0.14	*
2009 Mar 26.24	3.26	P60	<i>i'</i>	2880.0	22.38 ± 0.19	*
2009 Mar 27.31	4.31	P60	<i>i'</i>	5400.0	> 22.60	*
2009 Mar 27.36	4.36	Nickel	<i>R</i>	15600.0	22.79 ± 0.18	7
2009 Mar 28.11	5.10	TLS Tautenberg	<i>R</i>	6600.0	22.83 ± 0.20	8
2009 Mar 28.25	5.28	P60	<i>i'</i>	4500.0	22.92 ± 0.14	*
2009 Mar 28.89	5.89	TLS Tautenberg	<i>R</i>	2400.0	23.29 ± 0.50	9
2009 Mar 29.00	5.99	Shajn	<i>R</i>	5460.0	22.86 ± 0.1	10
2009 Mar 29.32	6.33	NOT	<i>R</i>	1800.0	22.96 ± 0.06	11
2009 Mar 31.91	8.90	TLS Tautenberg	<i>R</i>	7200.0	23.80 ± 0.36	9
2009 Apr 6.40	14.41	Gemini-N/GMOS	<i>r'</i>	1200.0	23.94 ± 0.10	*
2009 Jul 29.96	129.97	Gemini-S/GMOS	<i>r'</i>	1800.0	24.95 ± 0.15	*

<sup>a</sup> UT at beginning of exposure.

<sup>b</sup> Time from midpoint of exposure to *Fermi*-GBM trigger.

<sup>c</sup> Reported magnitudes have not been corrected for Galactic extinction ( $E(B - V) = 0.025$  mag; Schlegel et al. 1998). Observations in the *R* band are referenced to Vega, while all other filters are reported on the AB magnitude system (Oke & Gunn 1983).

<sup>d</sup> \* – This work; 1 – Updike et al. (2009b); 2 – Wang et al. (2009); 3 – Kann et al. (2009b); 4 – Burenin et al. (2009); 5 – Perley et al. (2009a); 6 – Guidorzi et al. (2009); 7 – Perley (2009); 8 – Kann et al. (2009c); 9 – Kann et al. (2009a); 10 – Rumyantsev & Pozanenko (2009); 11 – de Ugarte Postigo et al. (2009).

TABLE 10  
OPTICAL/NIR OBSERVATIONS OF GRB 090328

Date <sup>a</sup> (UT)	Time Since Burst <sup>b</sup> (days)	Telescope/Instrument	Filter	Exposure Time (s)	Magnitude <sup>c</sup>	Reference <sup>d</sup>
2009 Mar 29.06	0.67	UVOT	<i>U</i>	793.1	$19.38 \pm 0.10$	*
2009 Mar 29.13	0.73	UVOT	<i>U</i>	462.1	$19.58 \pm 0.16$	*
2009 Mar 29.98	1.58	Gemini-S/GMOS	<i>i'</i>	150.0	$19.98 \pm 0.10$	*
2009 Mar 29.98	1.60	GROND	<i>g'</i>	480.0	$20.97 \pm 0.05$	1
2009 Mar 29.98	1.60	GROND	<i>r'</i>	480.0	$20.23 \pm 0.03$	1
2009 Mar 29.98	1.60	GROND	<i>i'</i>	480.0	$19.89 \pm 0.04$	1
2009 Mar 29.98	1.60	GROND	<i>z'</i>	480.0	$19.54 \pm 0.03$	1
2009 Mar 29.98	1.60	GROND	<i>J</i>	480.0	$19.54 \pm 0.06$	1
2009 Mar 29.98	1.60	GROND	<i>H</i>	480.0	$19.02 \pm 0.06$	1
2009 Mar 29.98	1.60	GROND	<i>K</i>	480.0	$18.52 \pm 0.08$	1
2009 Mar 30.06	1.69	UVOT	<i>U</i>	4111.2	$20.80 \pm 0.11$	*
2009 Mar 31.42	3.04	UVOT	<i>U</i>	4162.2	$21.96 \pm 0.20$	*
2009 Apr 3.79	6.40	UVOT	<i>U</i>	2094.3	$21.52 \pm 0.24$	*
2009 Apr 4.79	7.40	UVOT	<i>U</i>	2507.4	$21.69 \pm 0.23$	*
2009 Apr 9.68	12.33	UVOT	<i>U</i>	9493.6	$22.67 \pm 0.24$	*

<sup>a</sup> UT at beginning of exposure.

<sup>b</sup> Time from midpoint of exposure to *Fermi*-GBM trigger.

<sup>c</sup> Reported magnitudes have not been corrected for Galactic extinction ( $E(B - V) = 0.057$  mag; Schlegel et al. 1998). Observations in the *U* band are referenced to Vega, while all other filters are reported on the AB magnitude system (Oke & Gunn 1983).

<sup>d</sup> \* – This work; 1 – Updike et al. (2009a).

TABLE 11  
OPTICAL OBSERVATIONS OF GRB 090902B

Date <sup>a</sup> (UT)	Time Since Burst <sup>b</sup> (days)	Telescope/Instrument	Filter	Exposure Time (s)	Magnitude <sup>c</sup>	Reference <sup>d</sup>
2009 Sep 2.51	0.06	ROTSE-IIIa	<i>R</i>	890.0	$16.4 \pm 0.5$	1
2009 Sep 2.72	0.27	ROTSE-IIIc	<i>R</i>	846.0	$> 18.7$	1
2009 Sep 2.98	0.53	UVOT	<i>U</i>	1074.7	$20.31 \pm 0.15$	*
2009 Sep 3.08	0.64	UVOT	<i>U</i>	3612.9	$20.54 \pm 0.09$	*
2009 Sep 3.19	0.74	Nickel	<i>R</i>	3300.0	$20.60 \pm 0.10$	1
2009 Sep 3.24	0.79	Gemini-N/GMOS	<i>r'</i>	180.0	$20.68 \pm 0.15$	*
2009 Sep 3.24	0.82	UVOT	<i>U</i>	8756.0	$20.92 \pm 0.08$	*
2009 Sep 3.33	0.88	Liverpool	<i>R</i>	1800.0	$21.04 \pm 0.11$	1
2009 Sep 3.74	1.30	UVOT	<i>U</i>	3852.8	$22.06 \pm 0.26$	*
2009 Sep 3.87	1.42	WHT-LIRIS	<i>J</i>	1800.0	$19.99 \pm 0.15$	1
2009 Sep 3.89	1.44	Liverpool	<i>R</i>	1800.0	$21.40 \pm 0.10$	1
2009 Sep 3.92	1.47	WHT-LIRIS	<i>K</i>	2592.0	$18.92 \pm 0.20$	1
2009 Sep 3.99	1.54	GROND	<i>r'</i>	738.0	$21.54 \pm 0.05$	1
2009 Sep 4.00	1.55	NOT	<i>V</i>	900.0	$21.67 \pm 0.11$	1
2009 Sep 4.01	1.56	NOT	<i>R</i>	900.0	$21.40 \pm 0.11$	1
2009 Sep 4.02	1.57	NOT	<i>I</i>	900.0	$20.72 \pm 0.11$	1
2009 Sep 4.36	1.91	UKIRT-WFCAM	<i>J</i>	1080.0	$20.20 \pm 0.20$	1
2009 Sep 4.36	1.91	UKIRT-WFCAM	<i>K</i>	1080.0	$18.90 \pm 0.25$	1
2009 Sep 4.98	2.53	GROND	<i>r'</i>	738.0	$22.01 \pm 0.07$	1
2009 Sep 5.93	3.48	Liverpool	<i>R</i>	1800.0	$22.60 \pm 0.25$	1
2009 Sep 6.37	3.97	UVOT	<i>U</i>	10549.1	$22.25 \pm 0.21$	*
2009 Sep 8.98	6.53	GROND	<i>r'</i>	738.0	$23.07 \pm 0.17$	1
2009 Sep 10.37	7.97	UVOT	<i>U</i>	9969.0	$> 22.56$	*

<sup>a</sup> UT at beginning of exposure.

<sup>b</sup> Time from midpoint of exposure to *Fermi*-GBM trigger.

<sup>c</sup> Reported magnitudes have not been corrected for Galactic extinction ( $E(B - V) = 0.042$  mag; Schlegel et al. 1998). Observations in the *r'* filter are reported on the AB magnitude system (Oke & Gunn 1983), while all other filters are referenced to Vega.

<sup>d</sup> \* – This work; 1 – Pandey et al. (2010).

TABLE 12  
OPTICAL/NIR OBSERVATIONS OF GRB 090926A

Date <sup>a</sup> (UT)	Time Since Burst <sup>b</sup> (d)	Telescope/Instrument	Filter (s)	Exposure Time	Magnitude <sup>c</sup>
2009 Sep 26.72	0.5415	UVOT	<i>U</i>	358.0	18.33 ± 0.06
2009 Sep 26.72	0.5501	UVOT	<i>V</i>	358.0	18.34 ± 0.10
2009 Sep 26.78	0.5989	UVOT	<i>U</i>	675.0	18.50 ± 0.05
2009 Sep 26.79	0.6149	UVOT	<i>V</i>	675.0	18.91 ± 0.09
2009 Sep 26.84	0.6649	UVOT	<i>U</i>	704.5	18.71 ± 0.05
2009 Sep 26.85	0.6816	UVOT	<i>V</i>	703.3	19.09 ± 0.10
2009 Sep 26.90	0.7299	UVOT	<i>U</i>	526.0	18.84 ± 0.07
2009 Sep 26.96	0.7966	PROMPT-2	<i>V</i>	560.0	18.76 ± 0.22
2009 Sep 26.96	0.7969	PROMPT-4	<i>R</i>	640.0	18.69 ± 0.16
2009 Sep 26.96	0.7986	PROMPT-5	<i>I</i>	400.0	18.02 ± 0.19
2009 Sep 26.98	0.8062	UVOT	<i>U</i>	459.4	18.94 ± 0.08
2009 Sep 26.99	0.8172	UVOT	<i>V</i>	459.4	18.97 ± 0.12
2009 Sep 27.00	0.8250	PROMPT-5	<i>I</i>	720.0	18.12 ± 0.07
2009 Sep 27.00	0.8259	PROMPT-4	<i>R</i>	880.0	18.49 ± 0.05
2009 Sep 27.00	0.8314	PROMPT-2	<i>V</i>	1600.0	19.12 ± 0.10
2009 Sep 27.00	0.8372	PROMPT-5	<i>I</i>	880.0	18.06 ± 0.05
2009 Sep 27.00	0.8379	PROMPT-4	<i>R</i>	880.0	18.45 ± 0.04
2009 Sep 27.00	0.8553	PROMPT-4	<i>R</i>	800.0	18.41 ± 0.04
2009 Sep 27.00	0.8553	PROMPT-5	<i>I</i>	800.0	18.04 ± 0.05
2009 Sep 27.00	0.8583	SMARTS / ANDICAM	<i>I</i>	2160.0	18.17 ± 0.07
2009 Sep 27.00	0.8583	SMARTS / ANDICAM	<i>J</i>	1800.0	17.28 ± 0.07
2009 Sep 27.00	0.8603	PROMPT-2	<i>V</i>	1520.0	19.12 ± 0.09
2009 Sep 27.04	0.8654	PROMPT-3	<i>B</i>	880.0	19.56 ± 0.10
2009 Sep 27.04	0.8659	PROMPT-4	<i>R</i>	800.0	18.36 ± 0.04
2009 Sep 27.04	0.8660	PROMPT-5	<i>I</i>	800.0	17.97 ± 0.05
2009 Sep 27.04	0.8662	UVOT	<i>U</i>	689.8	18.80 ± 0.06
2009 Sep 27.04	0.8761	PROMPT-4	<i>R</i>	800.0	18.36 ± 0.04
2009 Sep 27.04	0.8764	PROMPT-5	<i>I</i>	640.0	17.95 ± 0.05
2009 Sep 27.04	0.8808	PROMPT-2	<i>V</i>	1520.0	19.19 ± 0.09
2009 Sep 27.04	0.8811	PROMPT-3	<i>B</i>	1520.0	19.57 ± 0.07
2009 Sep 27.05	0.8826	UVOT	<i>V</i>	689.7	18.86 ± 0.09
2009 Sep 27.04	0.8858	PROMPT-5	<i>I</i>	800.0	17.87 ± 0.04
2009 Sep 27.04	0.8864	PROMPT-4	<i>R</i>	800.0	18.31 ± 0.04
2009 Sep 27.04	0.8962	PROMPT-5	<i>I</i>	800.0	17.92 ± 0.05
2009 Sep 27.04	0.8967	PROMPT-4	<i>R</i>	800.0	18.29 ± 0.04
2009 Sep 27.04	0.9012	PROMPT-3	<i>B</i>	1600.0	19.56 ± 0.07
2009 Sep 27.04	0.9013	PROMPT-2	<i>V</i>	1520.0	19.02 ± 0.08
2009 Sep 27.08	0.9055	PROMPT-5	<i>I</i>	640.0	17.85 ± 0.05
2009 Sep 27.08	0.9065	PROMPT-4	<i>R</i>	720.0	18.34 ± 0.04
2009 Sep 27.08	0.9172	PROMPT-4	<i>R</i>	880.0	18.29 ± 0.04
2009 Sep 27.08	0.9172	PROMPT-5	<i>I</i>	880.0	17.72 ± 0.03
2009 Sep 27.08	0.9223	PROMPT-3	<i>B</i>	1680.0	19.38 ± 0.05
2009 Sep 27.08	0.9225	PROMPT-2	<i>V</i>	1600.0	18.95 ± 0.07
2009 Sep 27.08	0.9280	PROMPT-4	<i>R</i>	800.0	18.29 ± 0.04
2009 Sep 27.08	0.9284	PROMPT-5	<i>I</i>	560.0	17.85 ± 0.05
2009 Sep 27.10	0.9331	UVOT	<i>U</i>	691.7	18.64 ± 0.05
2009 Sep 27.08	0.9388	PROMPT-4	<i>R</i>	880.0	18.15 ± 0.03
2009 Sep 27.08	0.9388	PROMPT-5	<i>I</i>	880.0	17.76 ± 0.04
2009 Sep 27.08	0.9419	PROMPT-3	<i>B</i>	1360.0	19.56 ± 0.08
2009 Sep 27.08	0.9421	PROMPT-2	<i>V</i>	1280.0	18.85 ± 0.08
2009 Sep 27.13	0.9475	PROMPT-4	<i>R</i>	480.0	18.25 ± 0.05
2009 Sep 27.13	0.9477	PROMPT-5	<i>I</i>	480.0	17.82 ± 0.05
2009 Sep 27.12	0.9495	UVOT	<i>V</i>	691.7	18.76 ± 0.09
2009 Sep 27.17	0.9933	PROMPT-3	<i>B</i>	960.0	19.60 ± 0.07
2009 Sep 27.17	0.9933	PROMPT-5	<i>I</i>	960.0	17.82 ± 0.04
2009 Sep 27.17	0.9935	PROMPT-2	<i>V</i>	960.0	18.88 ± 0.10
2009 Sep 27.17	0.9938	PROMPT-4	<i>R</i>	1040.0	18.21 ± 0.03
2009 Sep 27.17	0.9992	UVOT	<i>U</i>	554.0	18.65 ± 0.06
2009 Sep 27.17	1.0058	PROMPT-5	<i>I</i>	960.0	17.83 ± 0.03
2009 Sep 27.17	1.0062	PROMPT-4	<i>R</i>	880.0	18.24 ± 0.03
2009 Sep 27.17	1.0100	PROMPT-3	<i>B</i>	1200.0	19.43 ± 0.06
2009 Sep 27.17	1.0107	PROMPT-2	<i>V</i>	1520.0	19.03 ± 0.09
2009 Sep 27.17	1.0166	PROMPT-4	<i>R</i>	560.0	18.19 ± 0.04
2009 Sep 27.17	1.0173	PROMPT-5	<i>I</i>	640.0	17.79 ± 0.04
2009 Sep 27.21	1.0293	PROMPT-4	<i>R</i>	960.0	18.26 ± 0.03
2009 Sep 27.21	1.0295	PROMPT-5	<i>I</i>	880.0	17.85 ± 0.04
2009 Sep 27.21	1.0352	PROMPT-2	<i>V</i>	1680.0	18.89 ± 0.07
2009 Sep 27.21	1.0363	PROMPT-3	<i>B</i>	1360.0	19.53 ± 0.07
2009 Sep 27.21	1.0408	PROMPT-5	<i>I</i>	800.0	17.87 ± 0.04
2009 Sep 27.21	1.0412	PROMPT-4	<i>R</i>	880.0	18.43 ± 0.04
2009 Sep 27.21	1.0530	PROMPT-5	<i>I</i>	960.0	18.01 ± 0.04
2009 Sep 27.21	1.0530	PROMPT-4	<i>R</i>	960.0	18.42 ± 0.04
2009 Sep 27.21	1.0590	PROMPT-2	<i>V</i>	1840.0	19.13 ± 0.10
2009 Sep 27.21	1.0592	PROMPT-3	<i>B</i>	1920.0	19.64 ± 0.06

TABLE 12 — *Continued*

Date <sup>a</sup> (UT)	Time Since Burst <sup>b</sup> (d)	Telescope/Instrument	Filter (s)	Exposure Time	Magnitude <sup>c</sup>
2009 Sep 27.21	1.0653	PROMPT-4	<i>R</i>	960.0	18.43 ± 0.04
2009 Sep 27.21	1.0655	PROMPT-5	<i>I</i>	960.0	18.00 ± 0.05
2009 Sep 27.24	1.0658	UVOT	<i>U</i>	730.1	18.67 ± 0.05
2009 Sep 27.25	1.0777	PROMPT-4	<i>R</i>	960.0	18.56 ± 0.04
2009 Sep 27.25	1.0780	PROMPT-5	<i>I</i>	960.0	18.02 ± 0.04
2009 Sep 27.25	1.0839	PROMPT-3	<i>B</i>	1920.0	19.59 ± 0.05
2009 Sep 27.25	1.0839	PROMPT-2	<i>V</i>	1840.0	19.15 ± 0.09
2009 Sep 27.25	1.0900	PROMPT-4	<i>R</i>	960.0	18.45 ± 0.03
2009 Sep 27.25	1.0902	PROMPT-5	<i>I</i>	880.0	17.97 ± 0.04
2009 Sep 27.25	1.1028	PROMPT-5	<i>I</i>	1040.0	17.98 ± 0.04
2009 Sep 27.25	1.1029	PROMPT-4	<i>R</i>	1040.0	18.39 ± 0.03
2009 Sep 27.25	1.1091	PROMPT-3	<i>B</i>	2000.0	19.51 ± 0.04
2009 Sep 27.25	1.1093	PROMPT-2	<i>V</i>	1920.0	18.96 ± 0.07
2009 Sep 27.29	1.1158	PROMPT-4	<i>R</i>	960.0	18.47 ± 0.03
2009 Sep 27.29	1.1158	PROMPT-5	<i>I</i>	960.0	18.04 ± 0.04
2009 Sep 27.29	1.1283	PROMPT-5	<i>I</i>	960.0	18.02 ± 0.04
2009 Sep 27.29	1.1287	PROMPT-4	<i>R</i>	1040.0	18.45 ± 0.03
2009 Sep 27.29	1.1348	PROMPT-3	<i>B</i>	2000.0	19.62 ± 0.05
2009 Sep 27.29	1.1352	PROMPT-2	<i>V</i>	1920.0	19.09 ± 0.09
2009 Sep 27.29	1.1413	PROMPT-5	<i>I</i>	1040.0	17.97 ± 0.04
2009 Sep 27.29	1.1415	PROMPT-4	<i>R</i>	960.0	18.43 ± 0.03
2009 Sep 27.33	1.1545	PROMPT-4	<i>R</i>	960.0	18.47 ± 0.03
2009 Sep 27.33	1.1550	PROMPT-5	<i>I</i>	1040.0	18.06 ± 0.05
2009 Sep 27.33	1.1617	PROMPT-2	<i>V</i>	2000.0	19.11 ± 0.07
2009 Sep 27.33	1.1617	PROMPT-3	<i>B</i>	2080.0	19.67 ± 0.06
2009 Sep 27.33	1.1684	PROMPT-4	<i>R</i>	1040.0	18.51 ± 0.03
2009 Sep 27.33	1.1685	PROMPT-5	<i>I</i>	1040.0	18.06 ± 0.05
2009 Sep 27.33	1.1818	PROMPT-4	<i>R</i>	1040.0	18.52 ± 0.04
2009 Sep 27.33	1.1826	PROMPT-5	<i>I</i>	960.0	18.24 ± 0.06
2009 Sep 27.33	1.1850	PROMPT-3	<i>B</i>	1520.0	19.74 ± 0.08
2009 Sep 27.33	1.1859	PROMPT-2	<i>V</i>	1600.0	19.05 ± 0.10
2009 Sep 27.33	1.1919	PROMPT-4	<i>R</i>	480.0	18.63 ± 0.07
2009 Sep 27.33	1.1923	PROMPT-5	<i>I</i>	480.0	18.28 ± 0.11
2009 Sep 27.38	1.2093	PROMPT-3	<i>B</i>	480.0	19.73 ± 0.19
2009 Sep 27.38	1.2093	PROMPT-4	<i>R</i>	480.0	18.71 ± 0.08
2009 Sep 27.38	1.2093	PROMPT-5	<i>I</i>	480.0	18.18 ± 0.10
2009 Sep 27.38	1.2095	PROMPT-2	<i>V</i>	480.0	18.97 ± 0.23
2009 Sep 27.84	1.6702	UVOT	<i>U</i>	663.2	19.67 ± 0.10
2009 Sep 27.91	1.7349	UVOT	<i>U</i>	735.0	19.74 ± 0.09
2009 Sep 27.95	1.7846	UVOT	<i>V</i>	2168.5	19.82 ± 0.09
2009 Sep 27.97	1.8018	UVOT	<i>U</i>	737.0	19.71 ± 0.09
2009 Sep 28.04	1.8685	UVOT	<i>U</i>	742.9	19.78 ± 0.10
2009 Sep 28.11	1.9344	UVOT	<i>U</i>	577.3	19.79 ± 0.11
2009 Sep 28.12	1.9476	UVOT	<i>U</i>	341.3	19.73 ± 0.14
2009 Sep 28.40	2.2343	UVOT	<i>U</i>	1329.2	20.13 ± 0.08
2009 Sep 28.55	2.3829	UVOT	<i>U</i>	1730.2	20.01 ± 0.07
2009 Sep 28.56	2.3950	UVOT	<i>V</i>	1131.6	20.07 ± 0.14
2009 Sep 28.88	2.7199	UVOT	<i>U</i>	2642.7	20.11 ± 0.06
2009 Sep 28.89	2.7277	UVOT	<i>V</i>	2417.0	20.11 ± 0.10
2009 Sep 29.08	2.9301	PROMPT-5	<i>I</i>	3200.0	19.48 ± 0.10
2009 Sep 29.08	2.9346	PROMPT-4	<i>R</i>	2560.0	19.96 ± 0.12
2009 Sep 29.13	2.9642	PROMPT-5	<i>I</i>	1840.0	19.39 ± 0.11
2009 Sep 29.13	2.9644	PROMPT-4	<i>R</i>	1840.0	19.88 ± 0.12
2009 Sep 29.16	2.9892	SMARTS / ANDICAM	<i>I</i>	2160.0	19.53 ± 0.07
2009 Sep 29.16	2.9892	SMARTS / ANDICAM	<i>J</i>	1800.0	18.35 ± 0.09
2009 Sep 29.17	3.0041	PROMPT-5	<i>I</i>	2720.0	19.47 ± 0.10
2009 Sep 29.17	3.0055	PROMPT-3	<i>B</i>	3520.0	> 21.85
2009 Sep 29.17	3.0056	PROMPT-4	<i>R</i>	2960.0	19.85 ± 0.09
2009 Sep 29.17	3.0083	PROMPT-2	<i>V</i>	3600.0	> 20.94
2009 Sep 29.53	3.3664	UVOT	<i>V</i>	2533.9	20.75 ± 0.13
2009 Sep 29.54	3.3796	UVOT	<i>U</i>	3233.8	20.58 ± 0.07
2009 Sep 30.00	3.8439	PROMPT-5	<i>I</i>	5840.0	19.78 ± 0.11
2009 Sep 30.00	3.8446	PROMPT-4	<i>R</i>	6560.0	20.28 ± 0.10
2009 Sep 30.08	3.9295	PROMPT-5	<i>I</i>	6000.0	19.84 ± 0.12
2009 Sep 30.08	3.9302	PROMPT-4	<i>R</i>	5680.0	20.54 ± 0.14
2009 Sep 30.17	4.0116	PROMPT-3	<i>B</i>	30640.0	21.74 ± 0.13
2009 Sep 30.17	4.0121	PROMPT-2	<i>V</i>	28720.0	20.98 ± 0.17
2009 Sep 30.17	4.0249	PROMPT-5	<i>I</i>	7440.0	19.99 ± 0.12
2009 Sep 30.17	4.0253	PROMPT-4	<i>R</i>	7520.0	20.44 ± 0.09
2009 Sep 30.29	4.1428	PROMPT-5	<i>I</i>	10400.0	20.07 ± 0.11
2009 Sep 30.29	4.1445	PROMPT-4	<i>R</i>	10400.0	20.62 ± 0.10
2009 Sep 30.55	4.3878	UVOT	<i>V</i>	3091.7	20.60 ± 0.11
2009 Sep 30.55	4.3944	UVOT	<i>U</i>	3341.7	20.90 ± 0.08
2009 Oct 1.04	4.8842	PROMPT-4	<i>R</i>	12640.0	20.62 ± 0.10
2009 Oct 1.04	4.8854	PROMPT-5	<i>I</i>	12400.0	20.06 ± 0.10



TABLE 12 — *Continued*

Date <sup>a</sup> (UT)	Time Since Burst <sup>b</sup> (d)	Telescope/Instrument	Filter (s)	Exposure Time	Magnitude <sup>c</sup>
2009 Oct 1.11	4.9658	SMARTS / ANDICAM	<i>I</i>	2160.0	20.33 ± 0.10
2009 Oct 1.11	4.9658	SMARTS / ANDICAM	<i>J</i>	1800.0	19.16 ± 0.13
2009 Oct 1.13	4.9841	PROMPT-2	<i>V</i>	23520.0	> 21.84
2009 Oct 1.17	4.9896	PROMPT-3	<i>B</i>	24240.0	21.77 ± 0.17
2009 Oct 1.25	5.1003	PROMPT-5	<i>I</i>	12640.0	20.06 ± 0.11
2009 Oct 1.25	5.1024	PROMPT-4	<i>R</i>	11920.0	20.81 ± 0.13
2009 Oct 1.64	5.4915	UVOT	<i>U</i>	4142.8	21.36 ± 0.10
2009 Oct 2.04	5.8878	PROMPT-4	<i>R</i>	11600.0	20.72 ± 0.11
2009 Oct 2.17	5.9904	PROMPT-2	<i>V</i>	21360.0	> 21.74
2009 Oct 2.17	6.0045	PROMPT-3	<i>B</i>	28400.0	> 22.75
2009 Oct 2.25	6.0812	PROMPT-4	<i>R</i>	16960.0	20.93 ± 0.11
2009 Oct 2.25	6.0880	PROMPT-5	<i>I</i>	15040.0	20.50 ± 0.15
2009 Oct 3.00	6.8605	PROMPT-4	<i>R</i>	4240.0	> 21.23
2009 Oct 3.04	6.8617	PROMPT-5	<i>I</i>	4000.0	> 20.54
2009 Oct 3.04	6.8650	PROMPT-3	<i>B</i>	3520.0	> 21.45
2009 Oct 3.00	6.8781	UVOT	<i>U</i>	8789.3	21.59 ± 0.08
2009 Oct 4.13	7.9690	PROMPT-5	<i>I</i>	320.0	> 17.84
2009 Oct 4.13	7.9692	PROMPT-4	<i>R</i>	320.0	> 18.53
2009 Oct 4.13	7.9692	PROMPT-3	<i>B</i>	320.0	> 19.05
2009 Sep 4.74	8.5887	UVOT	<i>U</i>	4052.7	22.09 ± 0.18
2009 Oct 5.17	8.9975	PROMPT-4	<i>R</i>	12400.0	> 21.73
2009 Oct 5.17	9.0014	PROMPT-5	<i>I</i>	13280.0	> 21.24
2009 Oct 5.49	9.3449	UVOT	<i>U</i>	4640.7	22.07 ± 0.16
2009 Oct 6.38	10.2257	UVOT	<i>U</i>	4392.5	22.84 ± 0.29
2009 Oct 7.17	11.0096	PROMPT-4	<i>R</i>	27840.0	21.98 ± 0.22
2009 Oct 7.21	11.0377	PROMPT-5	<i>I</i>	23680.0	> 22.04
2009 Oct 7.42	11.2777	UVOT	<i>U</i>	5096.2	22.22 ± 0.17
2009 Oct 8.32	12.1742	UVOT	<i>U</i>	4573.2	22.94 ± 0.33
2009 Oct 19.05	22.90	Gemini-S / GMOS	<i>g'</i>	600.0	24.56 ± 0.11
2009 Oct 19.05	22.90	Gemini-S / GMOS	<i>r'</i>	600.0	23.97 ± 0.11
2009 Oct 19.05	22.90	Gemini-S / GMOS	<i>i'</i>	600.0	23.67 ± 0.12

<sup>a</sup> UT at beginning of exposure.<sup>b</sup> Time from midpoint of exposure to *Fermi*-GBM trigger.<sup>c</sup> Reported magnitudes have not been corrected for Galactic extinction ( $E(B - V) = 0.024$  mag; Schlegel et al. 1998). Observations in the  $g'$ -,  $r'$ -, and  $i'$ -bands are reported on the AB magnitude system (Oke & Gunn 1983). All other filters are referenced to Vega.